



# Smoking Prevalence Tracker 2020 Info-graph

(This information relates to data collected in the months of March, June, September and December 2020)

Survey carried out by Ipsos MRBI on behalf of the HSE

#### **Headline Data**

Smoking Prevalence: 15.4%

E Cigarette Use: 5.8%

Roll Your Own (RYO): 29.6%

(% of Smokers using RYO)



## Smoking Prevalence Tracker - Methodology

#### **Tracker Commenced in July 2002**

(Conducted monthly up to March 2016, now quarterly; results relate to surveys conducted in March, June, Sept. & December 2020)

Conducted quarterly via Omnipoll, Ipsos MRBI's telephone omnibus

(Conducted monthly up to March 2016)

N=1,000 adults 15+ per month Questionnaire covers smoking prevalence, brand & no. of cigarettes smoked & E-cigarette usage

Data provided in SPSS format for HSE analysis & reporting






### Differences in HSE Tracker and Healthy Ireland Smoking Prevalence Surveys

- The tracker survey is a smaller survey: (n=4000) taken quarterly whereas Healthy Ireland (HI) samples over 7,500 people annually.
- The tracker is a telephone survey whereas HI is a face to face household survey.
- The questions on smoking prevalence are slightly different (smoke one or more weekly in tracker whereas daily or occasionally in the Healthy Ireland survey).
- The tracker has been undertaken since 2002 whereas the Healthy Ireland survey started in 2015.
- Both surveys indicate an underlying downward trend in smoking prevalence, as indicated in Figure 1 for the tracker below:



Figure 1: Trend in smoking prevalence to 2020




Source: HSE Tracker Survey



## Cigarette Smoking Prevalence

Q.1 Do you smoke one or more cigarettes each week, whether packaged or roll your own?






# Cigarette Smoking Prevalence by Socio-Economic Group

Q.1 Do you smoke one or more cigarettes each week, whether packaged or roll your own?





Smoking prevalence is highest amongst those in the DE and C2 socio-economic groups, with those in the F & AB categories being least likely to smoke.

Social Class Categorisation Source: AIMRO Standard Guide for Social Class

- Further details available on HSE website

Base: All Respondents 2020: 4000