



Gemcitabine (1000mg/m²) and Capecitabine (830mg/m²) Therapy - 28 day

INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement status
Adjuvant treatment of pancreatic adenocarcinoma after	C25	00524a	Gemcitabine – Hospital
macroscopic complete resection			Capecitabine – CDS

TREATMENT:

The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patients individual clinical circumstances.

Gemcitabine is administered on day 1, 8 and 15 and capecitabine is taken on day 1-21 of a 28 day cycle for up to 6 cycles or until disease progression or unacceptable toxicity develops.

Facilities to treat anaphylaxis MUST be present when the chemotherapy is administered.

Day	Drug	Dose	Route	Diluent & Rate	Cycle
1,8,15	Gemcitabine	1000mg/m ²	IV infusion	250ml NaCl 0.9% over 30mins	Every 28 days
1 -21	Capecitabine	830mg/m ² Twice Daily ^a , ^{b, c}	РО	N/A	Every 28 days

 $^{{}^{\}mathrm{a}}\mathsf{The}$ dose to be administered should consider the available tablet strengths.

Reference the NCCP DOSE BANDING TABLES here for guidance on dosing of capecitabine.

Tablets should be swallowed whole with plenty of water with food or within 30 minutes of eating. Tablets should not be crushed or cut. b (Total daily dose = 1660mg/m²)

ELIGIBILITY:

- Indications as above
- ECOG 0-2

EXCLUSIONS:

- Hypersensitivity to gemcitabine, capecitabine or any of the excipients
- Known complete DPD deficiency
- History of severe and unexpected reactions to fluoropyrimidine therapy
- Pregnancy and lactation
- Severe hepatic or renal impairment

PRESCRIPTIVE AUTHORITY:

The treatment plan must be initiated by a Consultant Medical Oncologist.

NCCP Regimen: Gemcitabine(1000mg/m²) and Capecitabine (830mg/m²) Therapy-28 day	Published: 15/11/2018 Review: 23/09/2025	Version number: 4
Tumour Group: Gastrointestinal NCCP Regimen Code: 00524	ISMO Contributor: Prof Maccon Keane	Page 1 of 7

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer

^c See dose modifications section for patients with identified partial DPD deficiency.





TESTS:

Baseline tests:

- FBC, renal and liver profile
- DPD testing prior to first treatment with capecitabine using phenotype and/or genotype testing unless patient has been previously tested

Regular tests:

• Day 1: FBC, renal and liver profile

Day 8 and Day 15: FBC

Disease monitoring:

Disease monitoring should be in line with the patient's treatment plan and any other test/s as directed by the supervising Consultant.

DOSE MODIFICATIONS:

- Consider a reduced starting dose in patients with identified partial DPD deficiency.
 - Initial dose reduction may impact the efficacy of treatment. In the absence of serious toxicity, subsequent doses may be increased with careful monitoring.
- Any dose modification should be discussed with a Consultant.

Haematological:

Prior to commencing a new treatment cycle (i.e. day 1), ANC must be $\ge 1 \times 10^9 / L$ and platelets $\ge 100 \times 10^9 / L$.

Table 1: Dose modifications for gemcitabine and capecitabine within a cycle

ANC (x 10 ⁹ /L)		Platelet count (x 10 ⁹ /L)		Other toxicity	Recommended dose of Gemcitabine	Recommended dose of Capecitabine
≥1	and	≥100			100 %	100%
0.5- 1	or	50-100			75%	100%
< 0.5	or	<50			Omit. Do not restart treatment until ANC ≥ 0.5 and platelets ≥ 50	Interrupt treatment until recovery of toxicity to ≤ Grade 1
ANC < 0.5 for ≥ 5 days or ANC < 0.1 for ≥ 3 days or Any incidence of febrile neutropenia	or	< 25	or	cycle delay of >1 week due to any toxicity	Reduce dose to 75% of the original cycle initiation dose for all subsequent cycles.	Interrupt treatment until recovery of toxicity to ≤ Grade 1

NCCP Regimen: Gemcitabine(1000mg/m²) and Capecitabine (830mg/m²) Therapy-28 day	Published: 15/11/2018 Review: 23/09/2025	Version number: 4
Tumour Group: Gastrointestinal NCCP Regimen Code: 00524	ISMO Contributor: Prof Maccon Keane	Page 2 of 7

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer





Renal and hepatic impairment:

Table 2: Dose modifications for capecitabine and gemcitabine in renal and hepatic impairment

eatinine Clearance	D	
	Dose	In the absence of safety and efficacy
rCl ml/min)		data in patients with hepatic
≥30	100%	impairment, capecitabine use should be
<30	Omit	carefully monitored in patients with mild
		to moderate liver dysfunction,
		regardless of the presence or absence of
		liver metastasis.
≥30	100%	If bilirubin ≥27 micromol/L, use dose of
<30	Consider dose	800 mg/m ² and increase dose to full
	reduction – clinical	dose if tolerated.
	decision.	
	<30 ≥30 <30	<30 Omit ≥30 100% <30 Consider dose reduction – clinical

Management of adverse events:

Table 3: Dose Modification of gemcitabine for Adverse Events

Adverse reactions	Recommended dose modification of gemcitabine
Grade ≥ 2 Pneumonitis	Discontinue gemcitabine
Grade > 3 Non-haematological	Therapy with gemcitabine should be withheld (until toxicity has resolved
toxicity (except nausea/vomiting)	to grade ≤ 1) and may be resumed with 50% dose reduction or treatment
	discontinued at discretion of prescribing consultant.
Grade > 4 Non-haematological	Discontinue treatment
toxicity	

Table 4: Dose Modification of Capecitabine for Adverse Events

Toxicity grades*	Dose changes within a treatment cycle	Dose adjustment for next cycle/dose (% of starting dose)
Grade 1	Maintain dose level	Maintain dose level
Grade 2	Interrupt until resolved to grade 0-1	
 1st appearance 		100%
• 2 nd appearance		75%
3rd appearance		50%
• 4 th appearance	Discontinue permanently	
Grade 3	Interrupt until resolved to grade 0-1	
 1st appearance 		75%
• 2 nd appearance		50%
3rd appearance	Discontinue permanently	
Grade 4	Discontinue permanently OR	
• 1 st appearance	If consultant deems it to be in patient's best	50%
	interest to continue, interrupt until resolved	
	to grade 0-1	
• 2 nd appearance	Discontinue permanently	

Medication may be required for management of diarrhoea, e.g. loperamide (4mg at first onset followed by 2mg after each loose stool (max 16 mg /day) or see local policy.

"Common Terminology Criteria for Adverse Events (CTCAE) version 4	nology Criteria for Adverse Events (CTCAE) version 4.0.
---	---

NCCP Regimen: Gemcitabine(1000mg/m²) and Capecitabine (830mg/m²) Therapy-28 day	Published: 15/11/2018 Review: 23/09/2025	Version number: 4
Tumour Group: Gastrointestinal NCCP Regimen Code: 00524	ISMO Contributor: Prof Maccon Keane	Page 3 of 7

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer





Table 5: Dose Modification of capecitabine for diarrhoea

Grade	Diarrhoea	Dose changes within a treatment cycle	Dose adjustment for next cycle/dose (% of starting dose)
0-1	Increase of 2 to 3 stools/day or nocturnal stools	Maintain dose level	Maintain dose level
2	Increase of 4 to 6 stools/day or nocturnal stools		
	1 st appearance	Interrupt until resolved to grade 0-1	100%
	2 nd appearance		75%
	3rd appearance		50%
	4 th appearance	Discontinue permanently	
3	Increase of 7 to 9 stools/day or incontinence		
	1 st appearance	Interrupt until resolved to grade 0-1	75%
	2 nd appearance		50%
	3 rd appearance	Discontinue permanently	
4	Increase of 10 or more stools/day or grossly bloody diarrhoea; may require parenteral support		
	1 st appearance	Discontinue permanently or If consultant deems it to be in patient's best interest to continue, interrupt until resolved to grade 0-1	50%
	 2nd appearance 	Discontinue permanently	

Medication may be required for management of diarrhoea, e.g. loperamide (4mg at first onset followed by 2mg after each loose stool (max 16 mg/day) or see local policy

Hand foot syndrome:

Table 6: Dose modification of capecitabine in hand foot syndrome

Toxicity Grade		Dose Modification
Grade 1	Skin changes (e.g., numbness, dysesthesia, paraesthesia, tingling, erythema) with discomfort not disrupting normal activities	100% Dose
Grade 2	Skin changes (e.g., erythema, swelling) with pain affecting activities of daily living	Withhold treatment until event resolves or decreases in intensity to grade 1.
Grade 3	Severe skin changes (e.g., moist desquamation, ulceration, blistering) with pain, causing severe discomfort and inability to work or perform activities of daily living	Withhold treatment until event resolves or decreases in intensity to grade 1. Subsequent doses of capecitabine should be decreased

NCCP Regimen: Gemcitabine(1000mg/m²) and Capecitabine (830mg/m²) Therapy-28 day	Published: 15/11/2018 Review: 23/09/2025	Version number: 4
Tumour Group: Gastrointestinal NCCP Regimen Code: 00524	ISMO Contributor: Prof Maccon Keane	Page 4 of 7

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer
This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens





Treatment related hepatotoxicity

Table 7: Dose modification of capecitabine in treatment related hepatotoxicity

Bilirubin		ALT, AST	Dose Modification
> 3.0 x ULN	or	> 2.5 x ULN	Withhold treatment until bilirubin decreases to $\leq 3.0 \times ULN$ or ALT, AST decrease to $\leq 2.5 \times ULN$

SUPPORTIVE CARE:

EMETOGENIC POTENTIAL:

Gemcitabine: Low (Refer to local policy)

Capecitabine: Minimal to low (Refer to local policy)

PREMEDICATIONS: Not usually required

OTHER SUPPORTIVE CARE: Medication may be required for management of diarrhoea, e.g. loperamide (4mg at first onset followed by 2mg after each loose stool (max 16 mg /day) (Refer to local policy).

ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS

The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.

• **Neutropenia**: Fever or other evidence of infection must be assessed promptly and treated appropriately.

Gemcitabine:

- **Pulmonary Toxicity**: Acute shortness of breath may occur. Discontinue treatment with gemcitabine if drug-induced pneumonitis is suspected.
- Cardiovascular: Due to the risk of cardiac and/or vascular disorders with gemcitabine, particular caution must be exercised with patients presenting a history of cardiovascular events.
- **Haemolytic Uraemic syndrome:** Irreversible renal failure associated with hemolytic uremic syndrome may occur (rare) with gemcitabine. Use caution with pre-existing renal dysfunction.

Capecitabine:

- Diarrhoea and dehydration: This may be dose limiting with capecitabine therapy. Patients with severe
 diarrhoea should be carefully monitored and given fluid and electrolyte replacement if they become
 dehydrated.
- Cardiotoxicity: Angina-like chest pain, tachycardia, arrhythmias, heart failure, myocardial infarction
 and cardiac arrest may occur with capecitabine especially in patients with a prior history of coronary
 artery disease.
- **Dihydropyrimidine dehydrogenase (DPD) deficiency:** DPD is an enzyme encoded by the DPYD gene which is responsible for the breakdown of fluoropyrimidines. Patients with DPD deficiency are therefore at increased risk of fluoropyrimidine-related toxicity, including for example stomatitis, diarrhoea, mucosal inflammation, neutropenia and neurotoxicity. Treatment with 5-Fluorouracil,

NCCP Regimen: Gemcitabine(1000mg/m²) and Capecitabine (830mg/m²) Therapy-28 day	Published: 15/11/2018 Review: 23/09/2025	Version number: 4
Tumour Group: Gastrointestinal NCCP Regimen Code: 00524	ISMO Contributor: Prof Maccon Keane	Page 5 of 7

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer





capecitabine or tegafur-containing medicinal products is contraindicated in patients with known complete DPD deficiency. Consider a reduced starting dose in patients with identified partial DPD deficiency. Initial dose reduction may impact the efficacy of treatment. In the absence of serious toxicity, subsequent doses may be increased with careful monitoring. Therapeutic drug monitoring (TDM) of fluorouracil may improve clinical outcomes in patients receiving continuous 5-fluorouracil infusions.

• Hand-foot syndrome (HFS): HFS, also known as palmar-plantar erythrodysaesthesia (PPE), is a common side effect associated with capecitabine (see Table 6 for dose modification of capecitabine for HFS).

DRUG INTERACTIONS:

- Capecitabine enhances the anticoagulant effect of warfarin. Patients taking coumarin derivative anticoagulants should be monitored regularly for alterations in their coagulation parameters and the anti-coagulant dose adjusted accordingly.
- Sorivudine inhibits dihydropyrimidine dehydrogenase thus increasing its toxicity. Therefore capecitabine must not be administered concomitantly with sorivudine or its chemically related analogues.
- Patients taking phenytoin or fosphenytoin concomitantly with capecitabine should be regularly monitored for increased phenytoin plasma concentrations.
- Current drug interaction databases should be consulted for more information.

REFERENCES:

- 1. Neoptolemos J, Palmer D et al. Comparison of adjuvant Gemcitabine and Capecitabine with Gemcitabine monotherapy in patients with resected pancreatic cancer (ESPAC-4): a multicentre, open-label, randomised, phase 3 trial.
- 2. Knox J, Hedley D, Oza A. et al. Combining Gemcitabine and Capecitabine in Patients with Advanced Biliary Cancer: A phase 11 Trial. Journal of Clinical Oncology 2005;23(10), 2332-2338
- 3. BCCA Protocol Summary for Adjuvant Chemotherapy for Resected Pancreatic Adenocarcinoma Using Capecitabine and Gemcitabine- GIPAJGCAP (reviewed Nov 2017)
- 4. Dosage Adjustment for Cytotoxics in Renal Impairment January 2009; North London Cancer Network.
- 5. Dosage Adjustment for Cytotoxics in Hepatic Impairment January 2009; North London Cancer Network.
- 6. HPRA Direct Healthcare Professional Communication. 5-Fluorouracil (i.v.), capecitabine and tegafur containing products: Pre-treatment testing to identify DPD-deficient patients at increased risk of severe toxicity. Accessed Aug 2020 Available at: <a href="https://www.hpra.ie/docs/default-source/default-document-library/important-safety-information-from-marketing-authorisation-holders-of-products-containing-5-fluorouracil-(i-v-)-capecitabine-and-tegafur-as-approved-by-the-hpra.pdf?sfvrsn=0
- 7. NCCP Classification Document for Systemic Anti-Cancer Therapy (SACT) Induced Nausea and Vomiting. V4 2022. Available at:

 https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/nccp-classification-document-for-systemic-anti-cancer-therapy-sact-induced-nausea-and-vomiting.pdf
- Gemcitabine 40 mg/ml Concentrate for Solution for Infusion Summary of Product Characteristics.
 Accessed Sept 2020. Available at: https://www.hpra.ie/img/uploaded/swedocuments/Licence PA2059-039-001 18042019163629.pdf
- 9. Capecitabine (Xeloda®) Summary of Product Characteristics. Accessed Sept 2020. Available at:

NCCP Regimen: Gemcitabine(1000mg/m²) and Capecitabine (830mg/m²) Therapy-28 day	Published: 15/11/2018 Review: 23/09/2025	Version number: 4
Tumour Group: Gastrointestinal NCCP Regimen Code: 00524	ISMO Contributor: Prof Maccon Keane	Page 6 of 7

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer





http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-Product Information/human/000316/WC500058151.pdf

Version	Date	Amendment	Approved By
1	17/10/2018		Prof Maccon Keane
2	11/03/2020	Updated capecitabine dosing in renal impairment	Prof Maccon Keane
3	02/09/2020	Reviewed. Updated exclusion criteria, baseline testing, dose modifications and adverse events with respect to DPD deficiency as per DHPC from HPRA June 2020 Updated Adverse events regarding palmar-plantar erythrodysaesthesia.	Prof Maccon Keane
4	18/01/2023	Updated dose modification for gemcitabine in hepatic impairment. Amended emetogenic potential.	Prof Maccon Keane

Comments and feedback welcome at oncologydrugs@cancercontrol.ie.

NCCP Regimen: Gemcitabine(1000mg/m²) and Capecitabine (830mg/m²) Therapy-28 day	Published: 15/11/2018 Review: 23/09/2025	Version number: 4
Tumour Group: Gastrointestinal NCCP Regimen Code: 00524	ISMO Contributor: Prof Maccon Keane	Page 7 of 7

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer
This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens