

Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21-Day Therapy

INDICATIONS FOR USE:

| INDICATION | ICD10 | Regimen Code | Reimbursement Status |
|--|-------|--------------|----------------------|
| Treatment of newly diagnosed symptomatic multiple myeloma ⁱ | C90 | 00273a | Hospital |
| Treatment of relapsed/refractory multiple myeloma ⁱ | C90 | 00273b | Hospital |

TREATMENT:

The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patients individual clinical circumstances.

Treatment consists of four 3-week cycles of bortezomib administered on days 1, 4, 8, and 11; dexamethasone 40 mg on days 1, 4, 8, and 11; plus cyclophosphamide administered orally on days 1, 8, and 15 or until disease progression or unacceptable toxicity occurs.

*The dexamethasone dose may alternatively be administered as 20 mg on the day of and day after bortezomib (days 1, 2, 4, 5, 8, 9, 11 and 12) as per Kropff et al. 2007.

Table 1 Regimen Treatment Table

| Day | Drug | Dose | Route | Cycles |
|-------------|----------------------------|--|---------------------------------------|---------------------|
| 1, 4, 8, 11 | ^{a, b} Bortezomib | 1.3mg/m ² | ^{c, d} SC (abdomen or thigh) | 4 cycles of 21 days |
| 1, 8, 15 | Cyclophosphamide | ^e 300mg/m ² once daily | PO | 4 cycles of 21 days |
| 1, 4, 8, 11 | Dexamethasone | 40mg once daily | ^f PO | 4 cycles of 21 days |

At least 72 hours should elapse between consecutive doses of bortezomib.

^b Bortezomib is a proteasome inhibitor and is neurotoxic. Refer to [NCCP Guidance on the Safe Use of Neurotoxic drugs \(including Vinca Alkaloids\) in the treatment of cancer](#).

^c In individual cases where approved by Consultant, bortezomib may be administered as IV bolus over 3-5 seconds through a peripheral or central intravenous catheter followed by a flush with 0.9% NaCl. Note the concentration of bortezomib solution should be 1mg/ml when administered via the IV route.

^d The solution should be injected subcutaneously, at a 45-90° angle. Injection sites should be rotated for successive injections. If local injection site reactions occur, either a less concentrated solution may be administered SC or a switch to IV injection is recommended.

^e Cyclophosphamide is available as 50mg tablets. They should be swallowed with sufficient fluid without chewing. The tablets should not be divided before use. Higher doses up to 500mg/m² of cyclophosphamide may be used (Kumar et al).

^f Dexamethasone dose may alternatively be administered as 20 mg on the day of and day after bortezomib (days 1, 2, 4, 5, 8, 9, 11 and 12) as per Kropff et al. 2007.

Dose reduction of dexamethasone to 20mg or 10mg may be considered in selected patients depending on co morbidities. Dexamethasone to be taken in the morning with food.

| | | |
|--|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 1 of 8 |

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens

ELIGIBILITY:

- Indications as above
- ECOG 0-2
- Patients with pre-existing severe neuropathy should be treated with bortezomib only after careful risk/benefit assessment. Caution should be exercised as further treatment may result in severe prolonged neuropathy

EXCLUSIONS:

- Hypersensitivity to bortezomib, boron, dexamethasone or any of the excipients
- Acute diffuse infiltrative pulmonary and pericardial disease
- Pregnancy
- Lactation

PRESCRIPTIVE AUTHORITY:

The treatment plan must be initiated by a Consultant Haematologist working in the area of haematological malignancies

TESTS:

Baseline tests:

- FBC, renal, liver and bone profile
 - Blood pressure
 - Blood glucose* if being treated with oral hypoglycaemics (*See Drug Interactions)
 - Assessment of peripheral neuropathy status
 - Virology screen - Hepatitis B (HBsAg, HBcoreAb), hepatitis C, HIV
- *See Adverse Effects/Regimen Specific Complications re Hepatitis B Reactivation

Regular tests:

- FBC to be done minimum of day 1 and day 8 of each cycle
- Renal, liver and bone profile
- Blood pressure weekly
- Assessment of peripheral neuropathy status
- Blood glucose* if being treated with oral hypoglycaemics (*See Drug Interactions)

Disease monitoring:

Disease monitoring should be in line with the patient's treatment plan and any other test(s) as directed by the supervising Consultant.

| | | |
|---|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 2 of 8 |
| <p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p> | | |

DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Consultant.

Haematological toxicities:

Table 2: Dose Modification of Bortezomib and Cyclophosphamide for Haematological Toxicity

| Prior to Starting a new cycle | | |
|-------------------------------|---------------------------------|---|
| ANC (x10 ⁹ /L) | Platelets (x10 ⁹ /L) | Dose of Bortezomib and Cyclophosphamide |
| ≥0.5 | and ≥30 | 100% Dose |
| <0.5 | or <30 | Consider delay until recovery checking FBC weekly; reduce dose of bortezomib to 1mg/m ² . If the toxicity is not resolved or if it recurs at the lowest dose, discontinuation of bortezomib must be considered unless the benefit of treatment clearly outweighs the risk. |
| During a cycle | | |
| ANC (x10 ⁹ /L) | Platelets (x10 ⁹ /L) | Dose of Bortezomib and Cyclophosphamide |
| <0.5 | or <30 | Omit cyclophosphamide day 15. Withhold treatment with bortezomib until recovery of toxicity. Reinitiate treatment at a reduced dose of bortezomib (1.3 to 1mg/m ² or 1mg/m ² to 0.7mg/m ²) and consider dose reduction of cyclophosphamide. If the toxicity is not resolved or if it recurs at the lowest dose, discontinuation of bortezomib must be considered unless the benefit of treatment clearly outweighs the risk. |

Renal impairment:

Table 3: Dose Modification of Bortezomib and Cyclophosphamide in Renal Impairment

| Drug | Dose modification | |
|-------------------------|---|--------------------------|
| Bortezomib | It is unknown if the pharmacokinetics of bortezomib are influenced in patients with severe renal impairment not undergoing dialysis (CrCl < 20ml/min). Since dialysis may reduce bortezomib concentrations, it should be administered after the dialysis procedure. | |
| Cyclophosphamide | CrCl ml/min | Dose modification |
| | >20 | 100% |
| | 10-20 | 75% |
| | < 10 | 50% |
| | Clinical decision - consider whether patient is being treated with high dose treatment | |

| | | |
|---|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 3 of 8 |
| <p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p> | | |

Hepatic impairment:

Table 4: Dose Modification of Bortezomib and Cyclophosphamide in Hepatic Impairment

| Drug | Grade* | Bilirubin Level | SGOT (AST) levels | Modification of starting dose |
|---|--------------------------------------|-----------------|-------------------|---|
| Bortezomib | Mild | ≤1 x ULN | >ULN | None |
| | | >1 - 1.5 x ULN | Any | None |
| | Moderate | >1.5 - 3 x ULN | Any | Reduce bortezomib to 0.7 mg/m ² in the first treatment cycle. Consider dose escalation to 1.0 mg/m ² or further dose reduction to 0.5 mg/m ² in subsequent cycles based on patient tolerability. |
| | Severe | >3 x ULN | | |
| *Based on NCI Organ Dysfunction Working Group classification for categorising hepatic impairment (mild, moderate, severe). SGOT=serum glutamic oxaloacetic transaminase. | | | | |
| Cyclophosphamide | Severe impairment: Clinical decision | | | |

Neuropathic pain and/or peripheral neuropathy:

Table 5: Dose modifications for Bortezomib Related Neuropathy

| Severity of neuropathy | Dose Modification |
|--|---|
| Grade 1 (asymptomatic; loss of deep tendon reflexes or paresthesia) with no pain or loss of function | None |
| Grade 1 with pain or Grade 2 | Reduce dose to 1 mg/m ² <i>Or</i> Change treatment schedule to 1.3mg/m ² once every week |
| Grade 2 with pain or Grade 3 | Withhold treatment until symptoms of toxicity have resolved. When toxicity resolves re-initiate treatment and reduce dose to 0.7mg/m ² once every week. |
| Grade 4 and/or severe autonomic neuropathy | Discontinue treatment |
| Grade 1: Asymptomatic; clinical or diagnostic observations only Grade 2: Moderate symptoms; limiting instrumental Activities of Daily Living (ADL) Grade 3: Severe symptoms; limiting self-care ADL Grade 4: Life-threatening consequences; urgent intervention indicated <i>Grading based on NCI Common Toxicity Criteria CTCAE v 4</i> | |

| | | |
|---|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 4 of 8 |
| <p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p> | | |

Dose reductions for other toxicities:

Table 6: Dose Modification of Bortezomib for Adverse Events

| Adverse reactions | Recommended dose modification |
|--|---|
| Grade 3 Non-haematological toxicity | Withhold treatment until symptoms of the toxicity have resolved. Treatment may be reinitiated at a 25% reduced dose (1.3mg/m ² reduced to 1mg/m ² ; 1mg/m ² reduced to 0.7mg/m ²). If the toxicity is not resolved or if it recurs at the lowest dose, discontinuation of bortezomib must be considered unless the benefit of treatment clearly outweighs the risk. |
| New or worsening pulmonary symptoms (e.g. cough, dyspnoea) | Withhold treatment. Prompt diagnostic evaluation required and benefit/risk ratio should be considered prior to continuing bortezomib therapy. |
| Posterior Reversible Encephalopathy Syndrome (PRES) | Discontinue treatment. |

SUPPORTIVE CARE:

EMETOGENIC POTENTIAL:

Bortezomib: Low (**Refer to local policy**)
Cyclophosphamide: Moderate to high (**Refer to local policy**)

PREMEDICATIONS: Not usually required. Ensure patient remains well hydrated during treatment.

OTHER SUPPORTIVE CARE:

- Bisphosphonates should be considered in all patients with myeloma related bone disease.
- H₂-antagonist or PPI in patients receiving dexamethasone therapy (**Refer to local policy**).
- Low dose antiviral prophylaxis (**Refer to local policy**).
- Consider PJP prophylaxis (**Refer to local policy**).
- Tumour Lysis Syndrome prophylaxis (**Refer to local policy**)

ADVERSE EFFECTS/ REGIMEN SPECIFIC COMPLICATIONS:

The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.

Bortezomib:

- **Peripheral Neuropathy:** Patients with pre-existing severe neuropathy may be treated with bortezomib only after careful risk/benefit assessment.

| | | |
|---|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 5 of 8 |
| <p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p> | | |

- **Hypotension:** Treatment is commonly associated with orthostatic/postural hypotension. A minority of patients with orthostatic hypotension experienced syncopal events. Caution is advised when treating patients with a history of syncope receiving medicinal products known to be associated with hypotension; or who are dehydrated due to recurrent diarrhoea or vomiting.
- **Gastrointestinal toxicity:** Gastrointestinal toxicity, including nausea, diarrhoea, vomiting and constipation are very common with bortezomib treatment.
- **Hepatic Impairment:** Bortezomib is metabolised by liver enzymes. Bortezomib exposure is increased in patients with moderate or severe hepatic impairment; these patients should be treated with bortezomib at reduced doses and closely monitored for toxicities.
- **Haematological toxicity:** Gastrointestinal and intracerebral haemorrhage have been reported in association with bortezomib treatment. Therefore platelet counts should be monitored prior to each dose of bortezomib and bortezomib should be withheld when the platelet count is $<25 \times 10^9/L$. Potential benefit of treatment should be carefully weighed against the risks, particularly in case of moderate to severe thrombocytopenia and risk factors for bleeding. Complete blood counts with differential and including platelet counts should be frequently monitored throughout treatment with bortezomib. Platelet transfusion should be considered when clinically appropriate.
- **Seizures:** Seizures have been uncommonly reported in patients without previous history of seizures or epilepsy. Special care is required when treating patients with any risk factors for seizures.
- **Posterior Reversible Encephalopathy Syndrome (PRES):** In patients developing PRES, treatment with bortezomib should be discontinued.
- **Heart Failure:** Acute development or exacerbation of congestive heart failure, and/or new onset of decreased left ventricular ejection fraction has been reported during bortezomib treatment. Patients with risk factors for or existing heart disease should be closely monitored.
- **Renal Impairment:** Patients with renal impairment should be monitored closely.
- **Steroid use:** Steroid use is associated with numerous side effects including insomnia, gastric irritation, increased blood sugar levels, mood changes, increased appetite, bruising, skin fragility and osteoporosis (long term use).
- **Hepatitis B Reactivation:** Patients should be tested for both HBsAg and HBcoreAb as per local policy. If either test is positive, such patients should be treated with anti-viral therapy (Refer to local infectious disease policy). These patients should be considered for assessment by hepatology.

Cyclophosphamide

- **Haemorrhagic cystitis:** Ensure patient is well hydrated.

DRUG INTERACTIONS:

- Additive hypotensive effect with anti-hypertensives and bortezomib. Blood pressure should be monitored and ensure patient is well hydrated prior to bortezomib dose. Adjustment of anti-hypertensives may be required.
- During clinical trials, hypoglycemia was uncommonly reported and hyperglycemia commonly reported in diabetic patients receiving oral hypoglycemics. Patients on oral anti-diabetic agents receiving bortezomib treatment may require close monitoring of their blood glucose levels and adjustment of the dose of their anti-diabetic medications.

| | | |
|---|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 6 of 8 |
| <p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p> | | |

- Patients should be closely monitored when given bortezomib in combination with potent CYP3A4-inhibitors. Caution should be exercised when bortezomib is combined with CYP3A4- or CYP2C19 substrates. Current drug interaction databases should be consulted for more information.
- CYP3A-inhibitors also decrease the conversion of cyclophosphamide to both its active and inactive metabolites. Patients should also be counselled with regard to consumption of grapefruit juice.
- CYP3A-inducers increase the conversion of cyclophosphamide to both its active and inactive metabolites.
- Current drug interaction databases should be consulted for more information.

REFERENCES:

1. Reeder et al. Cyclophosphamide, bortezomib and dexamethasone (CyBorD) induction for newly diagnosed multiple myeloma: High response rates in a phase II clinical trial. *Leukaemia* 2009; 23(7): 1337–1341
2. Kropff M, Bisping G, Schuck, E. et al. Bortezomib in combination with intermediate-dose dexamethasone and continuous low-dose oral cyclophosphamide for relapsed multiple myeloma. *Br J Haematol* 2007; 138(3):330-337.
3. Kumar S et al. Randomized, multicenter, phase 2 study (EVOLUTION) of combinations of bortezomib, dexamethasone, cyclophosphamide, and lenalidomide in previously untreated multiple myeloma *Blood* 2012;119:4375-4382.
4. Moreau P, Coiteux V, Hulin C, et al. Prospective Comparison of subcutaneous versus intravenous administration of bortezomib in patients with multiple myeloma. *Haematologica* 2008; 93:1908-11.
5. Chanan-Kahn, Analysis of Herpes zoster events among bortezomib-treated patients. *J Clin Oncol.* 2008; 26:4784- 90.
6. Moreau P, Pylypenko H, Grosicki S, et al. Subcutaneous versus intravenous administration of bortezomib in patients with relapsed multiple myeloma: a randomized, phase 3, non-inferiority study. *Lancet Oncol.* 2011; 12(5):431- 40.
7. Dosage Adjustment for Cytotoxics in Renal Impairment January 2009; North London Cancer Network.
8. Dosage Adjustment for Cytotoxics in Hepatic Impairment January 2009; North London Cancer Network.
9. NCCP Classification Document for Systemic Anti-Cancer Therapy (SACT) Induced Nausea and Vomiting. V3 2021. Available at: <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/nccp-classification-document-for-systemic-anti-cancer-therapy-sact-induced-nausea-and-vomiting.pdf>
10. Bortezomib (VELCADE®) Summary of Product Characteristics, EMA. Last updated: 04/06/2021. Accessed July 2021. Available at: https://www.ema.europa.eu/en/documents/product-information/velcade-epar-product-information_en.pdf
11. Cyclophosphamide (Endoxana®) coated tablets 50mg. Summary of Product Characteristics. Accessed July 2021. Available at: https://www.hpra.ie/img/uploaded/swedocuments/Licence_PA2299-027-003_21122018112109.pdf

| | | |
|---|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 7 of 8 |
| <p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p> | | |

| Version | Date | Amendment | Approved By |
|---------|------------|--|------------------------------------|
| 1 | 05/04/2017 | | Dr Patrick Hayden Dr John Quinn |
| 2 | 19/06/2019 | Updated to new template. Updated recommendation on Hep B reactivation Updated dose modifications for haematological toxicity | Dr Patrick Hayden Dr John Quinn |
| 3 | 01/11/2021 | Reviewed. Amended treatment table. Updated exclusion criteria. Updated dose modification for hepatic impairment and for neuropathy. Updated emetogenic potential. Updated standard wording on Hep B reactivation. Updated adverse effects section to include gastrointestinal toxicity and haemorrhagic cystitis. | Dr Patrick Hayden Dr John Quinn |

Comments and feedback welcome at oncologydrugs@cancercontrol.ie.

ⁱ This is an unlicensed regimen for the use of bortezomib in Ireland. Patients should be informed of this and consented to treatment in line with the hospital's policy on the use of unlicensed medication and unlicensed or "off label" indications. Prescribers should be fully aware of their responsibility in communicating any relevant information to the patient and also ensuring that the unlicensed or "off label" indication has been acknowledged by the hospital's Drugs and Therapeutics Committee, or equivalent, in line with hospital policy

| | | |
|---|---|-------------------|
| NCCP Regimen: Cyclophosphamide, Bortezomib and Dexamethasone (CyBorD) 21 Day Therapy | Published: 05/04/2017 Review: 01/11/2026 | Version number: 3 |
| Tumour Group: Myeloma NCCP Regimen Code: 00273 | IHS/ISMO Contributor: Dr Patrick Hayden, Dr John Quinn | Page 8 of 8 |
| <p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p> | | |