

HSE 2018 PRESSURE ULCER CATEGORY / STAGING SYSTEM

Definition: “A pressure ulcer is a localised injury to the skin and / or underlying tissue usually over a bony prominence, as a result of pressure, or pressure in combination with shear. A number of contributing or confounding factors are also associated with pressure ulcers; the significance has yet to be elucidated”

Category / Stage I



Category/ Stage I: Intact skin with non – blanchable redness of a localised area usually over a bony prominence. Discolouration of the skin, warmth, odema, hardness or pain may also be present. Darkly pigmented skin may not have visible blanching. The area may be painful, firm, soft, warmer or cooler as compared to adjacent skin. (EPUAP 2009).

Category/Stage II



Category / Stage II: Partial thickness skin loss of dermis presenting as a shallow ulcer with a red pink wound bed, without slough. May present as an intact or open/ ruptured serum filled blister filled with serous or sero-sanguinous fluid. Presents as a shiny or dry shallow ulcer without slough or bruising. (EPUAP 2009).

Category/Stage III



Category / Stage III: Full thickness skin loss. Subcutaneous fat may be visible but bone, tendon or muscles are not exposed. Slough may be present but does not obscure the depth of tissue loss. The stage may include undermining or tunnelling (EPUAP 2009).

Category/Stage IV



Category / Stage IV: Full thickness tissue loss with exposed bone, tendon or muscle. Slough or eschar may be present. This stage often includes undermining and tunnelling. Exposed bone / muscle is visible or directly palpable (EPUAP 2009).

Suspected deep pressure and shear induced tissue damage, depth unknown



In individuals with non-blanchable redness and purple/maroon discoloration of intact skin combined with a history of prolonged, unrelieved pressure/shear, this skin change may be an indication of emerging, more severe pressure ulceration i.e. an emerging **Category/Stage III or IV Pressure Ulcer**. Clear recording of the exact nature of the visible skin changes, including recording of the risk that these changes may be an indication of emerging more severe pressure ulceration, should be documented in the patients' health record. These observations should be recorded in tandem with information pertaining to the patient history of prolonged, unrelieved pressure/shear. It is estimated that it could take **3-10 days** from the initial insult causing the damage, to become a **Category/Stage III or IV Pressure Ulcer** (Black et al, 2015).



Stable eschar (dry adherent, intact without erythema or fluctuance) on the heel serves as the body's biological cover and should not be removed. It should be documented as at least **Category / Stage III** until proven otherwise.