Forename:		Allergies / Adverse Drug Reactions						
	Medicine/Other	Nature of Reaction						
ATTENT LADEL ON								
DDRESSOGRAPH								
	Or No known allergies							
	Signature:	Date:						
	ATIENT LABEL OR DDRESSOGRAPH	Medicine/Other TIENT LABEL OR DDRESSOGRAPH Or No known allergies □						

Oxygen Therapy

Prescriber: For most chronic conditions, oxygen should be prescribed to achieve a target saturation of 94-98% (or 88-92% for those at risk of hypercapnic respiratory failure ie. CO ₂ retainers).										
Is the patient a known CO ₂ retainer ? Yes No Prescriber tick here if to continue upon discharge										
Prescription: Administration: Check and record flow rate (FR)/device (D) at each medicine round or other times specified and sign. Record oxygen saturations in the patient's observation chart										
Year: Day and Month ——— Other Times ———	>									
Continuous oxygen therapy□6Or 'When required' oxygen therapy□	1									
Target oxygen Saturation B8-92% 94-98% 10 Other saturation range FR	0									
Tick here □ if saturation not indicated and state reason D e.g. end of life care 12	2									
Starting device and flow rate:	4									
Prescriber Sig Reg No D Date 18	8									
Stop Date Reason Sig D										
Reviewed by Date FR	1									
Year: Day and Month Other Times										
Continuous oxygen therapy Or 'When required' oxygen therapy D	1									
Target oxygen Saturation □ 88-92% □ 94-98% 10 Other saturation range FR	D ?									
Tick here □ if saturation not indicated and state reason D e.g. end of life care 12	2									
Starting device and flow rate:	4									
Prescriber Sig Reg No D Date 18	8									
Stop Date Reason Sig D	2									
Reviewed by Date FR										

Device Codes

RA	Room Air (not requiring O2, weaning or on PRN O2)	CP Patient on CPAP system	V24 Venturi 24% (change figure as appropriate for % in use)			
SM	Simple mask	NV Patient on NIV System	H28 Humidified oxygen at 28% (change figure as			
Ν	Nasal Cannulae	OTH Other device	appropriate for percentage in use)			
RM	Reservoir mask	(specify)				
TM	Tracheostomy mask					
If a ward patient is requiring high flow ovviden via non represente mask, consider medical review						

If a ward patient is requiring high flow oxygen via non rebreathe mask, consider medical review. If target saturations are 88-92%, nebulised drugs should not be driven by oxygen (unless specified by the doctor).