

Step by Step guide - How to administer Intramuscular Naloxone

Further information on Prenoxad™ can be accessed through the below link http://www.prenoxadinjection.com/


Instructions for patient administration




Remove the clear film wrapping by pulling the tear strip on the side of the box. Twist the outer plastic box as shown to break the tamper evident seals and open.





The box contains 1 syringe of Prenoxad and two needles.







Peel back the backing paper from the needle packet and remove the needle in its protective sheath.








Gently twist the needle sheath and remove it from the syringe.



To inject someone who has overdosed, hold the syringe like a pen.



Insert the needle into the patient's outer thigh or upper arm, through clothing if necessary, and inject first dose (0.4ml). Withdraw the needle and syringe after each dose.

Step by Step guide - How to administer Intranasal Naloxone

Further information on Nyxoid™ can be accessed through the below link https://www.nyxoid.com/ie

 Peel off the back of the blister from the corner to remove the nasal spray from the container.

Place the nasal spray within easy reach.




2. Lay the person on their back.
Support the back of the neck and allow the head to tilt back.

Clear away anything blocking their nose.



3. Hold the nasal spray with your thumb on the bottom of the plunger and your first and middle fingers on either side of the nozzle.

Do not prime or test the Nyxoid™ nasal spray before use as it contains only one dose of naloxone and cannot be reused.



4. Gently insert the device nozzle in one nostril. Press firmly on the plunger until it <u>clicks</u> to give the dose.

Remove the nasal spray nozzle from the nostril after giving the dose.

