

Data Recording (continued)

Measurement 11	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 12	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 13	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 14	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 15	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 16	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 17	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 18	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 19	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	
Measurement 20	
Recording Date	
Weight	
Head Circumference	
Length/Height	
Location	
Health worker name	

GIRLS (Ireland) UK-WHO Growth Chart 0-4 years



Who should use this chart?
 Anyone who measures a child, plots or interprets charts should be suitably trained, or be supervised by someone qualified to do so. For further information and training materials see www.growthcharts.rcpch.ac.uk

A growth chart for all children
 The UK-WHO growth chart combines World Health Organization (WHO) standards with UK preterm and birth data. The chart from 2 weeks to 4 years of age is based on the WHO growth standard, derived from measurements of healthy, non-deprived, breastfed children of mothers who did not smoke.¹ The chart for birth measurements (32-42 weeks gestation) is based on British children measured around 1990.² The charts depict a healthy pattern of growth that is desirable for all children, whether breast fed or formula fed, and of whatever ethnic origin.³

Weighing and measuring

When measuring children up to 2 years, remove all clothes and nappy; children older than 2 years should wear minimal clothing only. Always remove shoes.

Weight: use only class III clinical electronic scales in metric setting.

Length: (before 2 years of age): proper equipment is essential (length board or mat). Measurers should be trained.

Height: (from 2 years): use a rigid rule with T piece, or stadiometer. Position head and feet as illustrated with child standing as straight as possible.

Head circumference: use a narrow plastic or paper tape to measure where the head circumference is greatest.

When to weigh
 When to weigh: Babies should be weighed in the first week as part of the assessment of feeding and thereafter as needed. Recovery of birthweight indicates that feeding is effective and that the child is well. Once feeding is established, babies should be weighed at routine healthchecks/immunisations. If there is concern weigh more often; however weights measured too close together are often misleading, so babies should be weighed no more than once a month up to 6 months of age, once every 2 months from 6 to 12 months of age, and once every 3 months over the age of 1 year. However, most children do not need to be weighed this often.

Please place sticker (if available) otherwise write in space provided.

Name: _____

Hospital/ID No:

Date of Birth:

When to measure length or height
 Length or height should be measured whenever there are any worries about a child's weight gain, growth or general health.

Plotting measurements
 For babies born at term (37 weeks or later), plot each measurement on the relevant chart by drawing a small dot where a vertical line through the child's age crosses a horizontal line through the measured value. The lettering on the charts ('weight', 'length' etc.) sits on the 50th centile, providing orientation for ease of plotting.

Plot birth weight (and, if measured, length and head circumference) at age 0 on the 0-1 year chart. The coloured arrows at age 0 represent UK birth weight data and show the child's birth centile.

Weight gain in the early days varies a lot from baby to baby, so there are no lines on the chart between 0 and 2 weeks. However, by 2 weeks of age most babies will be on a centile close to their birth centile.

For **preterm infants** a separate low birth weight chart is available for infants of less than 32 weeks gestation and any other infant requiring detailed assessment. For healthy infants born from 32 weeks and before 37 weeks, plot all measurements in the preterm section (to the left of the main 0-1 year chart) until 42 weeks gestation, then plot on the 0-1 year chart using gestational correction, as shown below.

The preterm section can also be used to assess the relative size of infants at the margin of 'term' (e.g. 37 weeks gestation), but these measurements should also be plotted at age 0 on the 0-1 year chart.

Gestational correction
 Plot measurements at the child's actual age and then draw a line back the number of weeks the infant was preterm. Mark the spot with an arrow (see diagram): this is the child's gestationally corrected centile. Gestational correction should continue until at least 1 year of age.

Centile terminology

If the point is within 1/4 of a space of the line they are on the centile: e.g. 91st.

If not they should be described as being between the two centiles: e.g. 75th-91st.

A centile space is the distance between two of the centile lines, or equivalent distance if midway between centiles.

Plotting for preterm infants (less than 37 weeks gestation):
 Draw a line back the number of weeks preterm and mark spot with arrow.

Interpreting the chart
Assessing weight loss after birth
 Most babies lose some weight after birth but 80% will have regained this by 2 weeks of age. Fewer than 5% of babies lose more than 10% of their weight at any stage; only 1 in 50 are 10% or more lighter than birth weight at 2 weeks.

Percentage weight loss can be calculated as follows:

$$\text{Percentage weight loss} = \frac{\text{Weight loss}}{\text{Birth weight}} \times 100\%$$

For example, a child born at 3.500kg who drops to 3.150kg at 5 days has lost 350g or 10%; in a baby born at 3.000kg, a 300g loss is 10%.

Careful clinical assessment and evaluation of feeding technique is indicated when weight loss exceeds 10% or recovery of birth weight is slow.

What do the centiles mean?
 These charts indicate a child's size compared with children of the same age and maturity who have shown optimum growth. The chart also shows how quickly a child is growing. The centile lines on the chart show the expected range of weights and heights (or lengths); each describes the number of children expected to be below that line (e.g. 50% below 50th, 91% below the 91st). Children come in all shapes and sizes, but 99 out of 100 children who are growing optimally will be between the two outer lines (0.4th and 99.6th centiles); half will lie between the 25th and 75th centile lines.

Being very small or very big can sometimes be associated with underlying illness. There is no single threshold below which a child's weight or height is definitely abnormal, but only 4 per 1000 children who are growing optimally are below the **0.4th centile**, so these children should be assessed at some point to exclude any problems. Those above the **99.6th centile** for height are almost always healthy. Also calculate BMI if weight and height centiles appear very different.

What is a normal rate of weight gain and growth?
 Babies do not all grow at the same rate, so a baby's weight often does not follow a particular centile line, especially in the first year. Weight is most likely to track within one centile space (the gap between two centile lines, see diagram). In infancy, acute illness can lead to sudden weight loss and a weight centile fall but on recovery the child's weight usually returns to its normal centile within 2-3 weeks. However, a sustained drop through two or more weight centile spaces is unusual (fewer than 2% of infants) and should be carefully assessed by the primary care team, including measuring length/height.


Because it is difficult to measure length and height accurately in pre-school children, successive measurements commonly show wide variation. If there are worries about growth, it is useful to measure on a few occasions over time; most healthy children will show a stable average position over time.

UK children have relatively large heads compared to the WHO standard, particularly after the age of 6 months. After the age of 6 weeks a head circumference below the 2nd centile will be seen in only 1 in 250 children. A head circumference above the 99.6th centile, or crossing upwards through 2 centile spaces, should only cause concern if there is a continued rise after 6 months, or other signs or symptoms.

Why do the length/height centiles change at 2 years?
 The growth standards show length data up to 2 years of age, and height from age 2 onwards. When a child is measured standing up, the spine is squashed a little, so their height is slightly less than their length; the centile lines shift down slightly at age 2 to allow for this. It is important that this difference does not worry parents; what matters is whether the child continues to follow the same centile after the transition.

Predicting adult height
 Parents like to know how tall their child will be as an adult. The child's most recent height centile (aged 2-4 years) gives a good idea of this for healthy children. Plot this centile on the adult height predictor to the right of the height chart to find the average adult height for children on this centile. Four out of five children will have adult heights that are within 6cm above or below this value.

Weight-height to BMI conversion chart
 BMI indicates how heavy a child is relative to his or her height and is the simplest measure of thinness and fatness from the age of 2, when height can be measured fairly accurately. This chart⁴ provides an approximate BMI centile, accurate to a quarter of a centile space.



Date:

Age:

BMI Centile:

- Instructions for use**
1. Read off the weight and height centiles from the growth chart.
 2. Plot the weight centile (left axis) against the height centile (bottom axis) on the chart above.
 3. If between centiles, read across in this position.
 4. Read off the corresponding BMI centile from the slanting lines.
 5. Record the centile with the date and child's age in the data box.

Interpretation
 In a child over 2 years of age, the BMI centile is a better indicator of overweight or underweight than the weight centile; a child whose weight is average for their height will have a BMI between the 25th and 75th centiles, whatever their height centile. BMI above the 91st centile suggests that the child is overweight; a child above the 98th centile is very overweight (clinically obese). BMI below the 2nd centile is unusual and may reflect undernutrition.

References

1. www.who.int/childgrowth/en
2. Cole TJ, Freeman JV, Preece MA. British 1990 growth reference centiles for weight, height, body mass index and head circumference fitted by maximum penalized likelihood. Stat Med 1998;17:407-29.
3. www.sacn.gov.uk/reports_position_statements/index.html
4. Cole TJ. A chart to link child centiles of body mass index, weight and height. Eur J Clin Nutr 2002;56:1194-9.