

Class: Date of Birth: Name: School Roll Number: Client ID:

4. Has your child been diagnosed with asthma? Yes No
5. If yes does your child take regular steroids for their asthma and/or has your child ever been admitted to ICU/Critical Care for Asthma? Yes No
6. Does your child take aspirin/salicylates medication? Yes No
7. Does your child have a severely weakened immune system due to disease or treatment? e.g., leukaemia/lymphoma or high dose steroids or severe neutropenia? Yes No
8. Does your child live with anyone currently having treatment that severely affects their immune system? e.g., someone who has had a bone marrow transplant? Yes No
9. Does your child take medication called combination checkpoint inhibitors e.g., ipilimumab plus nivolumab? Yes No
10. Is your child known to have a condition causing a Cerebrospinal Fluid (CSF) leak and/or has your child had a recent cochlear implant? Yes No
11. Did your child receive the flu vaccine last year? Yes No
12. Has your child ever received the flu vaccine before? Yes No
13. Does your child have severe neutropenia (low levels of a type of white blood cell) i.e. absolute neutrophil count $<0.5 \times 10^9 /L$? This does not apply to those with primary autoimmune neutropenia. Yes No

Some children aged under 9 years who have never had a flu vaccine before will need to have a second dose of flu vaccine if they are at high risk of complications from infection with flu. If your child has never had a flu vaccine before and is aged under 9 years, they may need a second dose of flu vaccine if they have any of the following conditions:

- Chronic heart disease
- Chronic liver disease
- Chronic neurological disease
- Chronic renal failure
- Chronic respiratory disease (including cystic fibrosis, moderate or severe asthma)
- Diabetes mellitus
- Any condition that might mean they cannot breathe well (e.g., a spinal cord injury, seizure disorder, or other neuromuscular disorder)
- Down syndrome
- Cancer
- Immunosuppression due to disease or treatment including asplenia or hyposplenism
- Moderate to severe neurodevelopmental disorders such as cerebral palsy and intellectual disability

Does your child have one or more of the conditions listed above? Yes No

Before your child is vaccinated you should let your vaccinator know if your child:

- has had influenza antiviral medications in the 48 hours before their vaccine is due, they should not get the vaccine.
- has an acute exacerbation of asthma, including increased wheezing and/or needed additional inhalers in the previous 72 hours they should not receive the nasal flu vaccine.
- has gotten a dose of the flu vaccine from their GP or Pharmacist since the consent form was completed, they should not get the vaccine.
- is unwell with a sudden fever (as vaccination should be delayed until recovery).

You can contact the vaccinator by phoning them.

Please note that this vaccine is not suitable for people who are pregnant.

