

Vaccine	4 months to <12 months	12 months to < 2 years	2 years to < 4 years	4 to <10 years	10 to <18 years	18 years and older
<b>6 in 1<sup>1</sup></b>	3 doses 2 months apart	3 doses 2 months apart	3 doses 2 months apart	3 doses 2 months apart		
<b>Men B</b>	2 doses 2 months apart (1 dose if ≥ 10 months and booster at 12 months or older, 2 months after the first dose)	2 doses 2 months apart <b>(if born on or after October 1<sup>st</sup> 2016)</b>	2 doses 1 month apart <b>(if born on or after October 1<sup>st</sup> 2016)</b>			
<b>Men C</b>	1 dose	1 dose	1 dose	1 dose	1 dose (if given at ≥ 10 years, adolescent booster not required)	1 dose (up to 23 years )
<b>PCV</b>	2 doses 2 months apart	1 dose (omit if ≥2 years of age <sup>2</sup> )				
<b>MMR<sup>3</sup></b>		1 dose	1 dose	2 doses 1 month apart	2 doses 1 month apart	2 doses 1 month apart <sup>4</sup>
<b>Rotavirus<sup>5</sup></b>	2 doses 4 weeks apart <b>No dose after 8 months 0 days</b>					
<b>Tdap/IPV</b>					3 doses 1 month apart	1 dose <sup>6</sup>
<b>Td/IPV</b>						2 doses - 1 month apart (1 month after Tdap/IPV)
<b>NOTE</b>	<i>Continue with routine childhood immunisation schedule from 12 months</i>		<i>Continue with routine school immunisations</i>  <i>Tdap/IPV at least 6 months and preferably 3 years after primary course MMR at least 1 month after previous</i>	<i>Continue with routine school immunisations</i> <i>Tdap/IPV at least 6 months and preferably 3 years after primary course]</i>	<i>Booster of Tdap/IPV 5 years after primary course and Tdap 10 years later</i>	

<sup>1</sup> One dose of single Hib vaccine may be given to children over 12 months of age and up to 10 years of age if this is the only vaccine they require

<sup>2</sup> Unless at increased risk

<sup>3</sup> 2<sup>nd</sup> dose of MMR is recommended routinely at 4-5 years but may be administered earlier. Children vaccinated <12 months in the case of an outbreak should have a repeat MMR vaccination at 12 months of age, at least one month after 1<sup>st</sup> vaccine with a further dose at 4-5 years of age. If a child aged <18 months receives a 2<sup>nd</sup> MMR vaccine within 3 months of the 1<sup>st</sup> MMR a 3<sup>rd</sup> MMR should be given at 4-5 years of age.

<sup>4</sup> For health care workers without presumptive evidence of immunity; for contacts in outbreaks born in Ireland or born outside Ireland and for adults from low resource countries, without evidence of two doses of MMR vaccine

<sup>5</sup> One dose if 7-<8 months

<sup>6</sup> Only one dose of Tdap/IPV is required due to likely previous exposure to pertussis infection