



# 4 in 1 and MMR School Vaccination Programme for Children in Junior Infants

Name: \_\_\_\_\_

Date: \_\_\_ / \_\_\_ / \_\_\_\_\_ Time of vaccination: \_\_\_ / \_\_\_



## Your child was given the following vaccines today

4 in 1



MMR



## Common reactions expected after these vaccines may include

- Mild fever.
- Soreness, swelling and redness where the injection was given. Sometimes swelling after 4 in 1 vaccine can be from the shoulder to the elbow. This swelling begins within 2 days of the vaccination and gets better over 4-5 days. Antibiotics are not needed to treat this local reaction.

## After MMR vaccine

- Some children may get “mini measles” with a rash and fever 6 to 10 days after the injection.
- On rare occasions, children may get “mini-mumps” with swelling in the jaw in the third week after vaccination.

These are not contagious.

## You can give your child paracetamol or ibuprofen to relieve aches and pains or to lower the fever.

If you are concerned about your child the school vaccination team can be contacted during office hours from Monday to Friday at

If you require medical advice after these hours please contact your GP. You will get a record of your Child’s Vaccinations when they have all been completed. **Please keep this safe.**

Your child was given the following vaccines today:

Tetravac  MMRVaxPro  Priorix

05/2024  
Order Code: HNI00876



For more information see: [www.immunisation.ie](http://www.immunisation.ie)

You can also find links to the patient information leaflets for the vaccines at [www.hpra.ie](http://www.hpra.ie). Search for Tetravac (4 in 1), Priorix or MMRVaxPro (MMR) to read the relevant patient information leaflet.