

# Immunisation Passport



Name

DOB

This is an important document,  
please keep it safe

[www.immunisation.ie](http://www.immunisation.ie)

For  
children born  
on or after  
1st July  
2015

## Introduction

The aim of this Immunisation Passport is to ensure you have a record of all your immunisations.

These records should include all

- primary childhood immunisations,
- school immunisations,
- travel, and
- adult immunisations

There are times when you may require other vaccines following an outbreak or for work or travel reasons. You can also record those vaccines in this booklet.

**Please bring this Immunisation Passport with you every time you attend an immunisation appointment.**



Name

Also known as

Date of Birth

Place of Birth

Address

Contact Phone Number(s)

Mobile

Email Address

Individual Health Identifier

PPS No.

Allergies

In Case of Emergency

Name(s)

Contact Number(s)

GP Name

GP Address

GP Phone Number

GP Name

GP Address

GP Phone Number

GP Name

GP Address

GP Phone Number

**Please keep this  
booklet safe and  
bring it with you to  
every vaccination  
appointment.**



The table below shows at what age the immunisations are given, where they are given and which vaccines are given.

**All the immunisations listed are free.**

When	Where	Vaccine(s)
At birth	HSE clinic or maternity hospital	BCG
2 months	GP surgery Visit 1	6 in 1 + PCV
4 months	GP surgery Visit 2	6 in 1 + MenC
6 months	GP surgery Visit 3	6 in 1 + PCV
12 months	GP surgery Visit 4	MMR + PCV
13 months	GP surgery Visit 5	MenC + Hib
Junior infants (first year of primary school)	School*	4 in 1 + MMR
First year of second level school (girls only)	School	HPV (2 doses)
First year of second level school	School	Tdap
First year of second level school	School	MenC
Every year (Between October and April)	GP Surgery or pharmacy	Flu**

\* These vaccines are only available in GP surgeries in Sligo, Leitrim and Donegal.

\*\* For those aged 65 and older and in at risk groups.

BCG	Bacille Calmette-Guérin (TB)
6 in 1	Diphtheria, Haemophilus influenzae b (Hib), Hepatitis B, Pertussis (Whooping cough), Polio, Tetanus
PCV	Pneumococcal conjugate
MenC	Meningococcal C
MMR	Measles, Mumps, Rubella
Hib	Haemophilus influenzae b
4 in 1	Diphtheria, Pertussis (whooping cough), Polio, Tetanus
HPV	Human papillomavirus
Tdap	Tetanus, low dose diphtheria and acellular pertussis (whooping cough)
Flu	Influenza

# Primary Childhood Immunisations

(for babies born on or after 1st July 2015)

Vaccine name	Age given	Manufacturer	Batch number
BCG			

## GP Visit 1 (2 months)

6 in 1			
PCV			

## GP Visit 2 (4 months)

NEXT APPOINTMENT DATE

6 in 1			
MenC			

## GP Visit 3 (6 months)

NEXT APPOINTMENT DATE

6 in 1			
PCV			

BCG Bacille Calmette-Guérin (TB)  
6 in 1 Diphtheria, Haemophilus influenzae b (Hib), Hepatitis B, Pertussis (Whooping cough), Polio, Tetanus



# Primary Childhood Immunisations

(for babies born on or after 1st July 2015)

Vaccine name	Age given	Manufacturer	Batch number
--------------	-----------	--------------	--------------

## GP Visit 4 (12 months)

NEXT APPOINTMENT DATE

MMR			
PCV			

## GP Visit 5 (13 months)

NEXT APPOINTMENT DATE

MenC			
Hib			

## Other


MMR Measles, Mumps, Rubella  
PCV Pneumococcal conjugate



Name

DOB

Expiry date	Route/site	Administered by	Date
-------------	------------	-----------------	------










**MenC** Meningococcal C  
**Hib** Haemophilus influenzae b

# School Immunisations

Vaccine name	Age given	Manufacturer	Batch number
4 in 1			
MMR			
Tdap			
MenC			
HPV Dose 1			
NEXT APPOINTMENT DATE			
HPV Dose 2			
Other			

- 4 in 1 Diphtheria, Pertussis (whooping cough), Polio, Tetanus  
MMR Measles, Mumps, Rubella  
Tdap Tetanus, low dose diphtheria and acellular pertussis (whooping cough)







# Post Vaccination Advice

**What common reactions can my child get after being vaccinated and what should I do?**

Common reaction	What to do
Soreness, swelling* and redness in the area where the injection was given	Give paracetamol or ibuprofen to relieve aches and pains.
	Make sure clothes are not too tight or rubbing against the area where the injection was given.
Fever (over 39.5°C)	Do not overdress your child.
	Make sure their room isn't too hot.
	Give extra fluids to drink.
	Give paracetamol or ibuprofen to lower the fever.
Headache or irritability	Give paracetamol or ibuprofen to relieve aches and pains.

\* In school children this swelling can be from the shoulder to the elbow. This usually occurs within 2 days of the vaccination and gets better over 4 – 5 days. Antibiotics are not needed to treat this local reaction.



- The dose of paracetamol or ibuprofen recommended for your child is written on the bottle according to the child's age.
- Please ask your pharmacist for sugar-free paracetamol or ibuprofen suitable for your child's age.
- Using paracetamol or ibuprofen over a long period without advice from a doctor may be harmful.



If your child received the MMR vaccine they may get a rash 6-10 days later (mini measles). This is not contagious.

## Notes

---

---

---

---

---

---

---

---

---

---

Published by: HSE National Immunisation Office  
Publication date: July 2015  
Order code: HNI00936