



4 in 1 and MMR School Vaccination Programme for Children in Junior Infants

Name: _____

Date: ____ / ____ / ____ Time of vaccination: ____ / ____



Your child was given the following vaccines today

4 in 1



LEFT
ARM



RIGHT
ARM

MMR



LEFT
ARM



RIGHT
ARM

Common reactions expected after these vaccines may include

- Mild fever.
- Soreness, swelling and redness where the injection was given. Sometimes swelling after 4 in 1 vaccine can be from the shoulder to the elbow. This swelling begins within 2 days of the vaccination and gets better over 4-5 days. Antibiotics are not needed to treat this local reaction.

After MMR vaccine

- Some children may get "mini measles" with a rash and fever 6 to 10 days after the injection.
- On rare occasions, children may get "mini-mumps" with swelling in the jaw in the third week after vaccination.

These are not contagious.

You can give your child paracetamol or ibuprofen to relieve aches and pains or to lower the fever.

If you are concerned about your child the school vaccination team can be contacted during office hours from Monday to Friday at

If you require medical advice after these hours please contact your GP.
You will get a record of your Child's Vaccinations when they have all been completed.
Please keep this safe.

Your child was given the following vaccines today:

Tetravac ☐ MMRVaxPro ☐ Priorix ☐

05/2024
Order Code: HNI00876

For more information see: www.immunisation.ie

You can also find links to the patient information leaflets for the vaccines at www.hpra.ie. Search for Tetravac (4 in 1), Priorix or MMRVaxPro (MMR) to read the relevant patient information leaflet.



Clár Vacsaínithe Scoile 4 in 1 agus Teanndáileog MMR do Pháistí sna Naíonáin Sóisearacha

Ainm: _____

Dáta: ____ / ____ / ____ Am an vacsaínithe: ____ / ____



Tugadh na vacsaíní seo a leanas dod' pháiste inniu

4 in 1



MMR



Cuimsítear sna frithghníomhuíthe coitianta a mbítear ag súil leo i ndiaidh na vacsaíní sin

- Fiabhras éadrom.
- Pian, at agus deirge san áit ar tugadh an t-instealladh. In amanna, d'fhéadfaí go mbeadh an t-at ón ngualainn go dtí an uillinn i ndiaidh na vacsaíne ceithre ghalar. Tosaíonn an t-at seo laistigh de dhá lá i ndiaidh an vacsaín a fháil agus feabhsaíonn rudaí thar thréimhse 4-5 lá. Níl gá le hantaibheathaigh chun an frithghníomhú logánta seo a chóireáil.

Tar éis an vacsaínithe MMR

- D'fhéadfadh roinnt pháistí an 'bhrúitíneach bheag' a tholg le gríos craicinn agus fiabhras 6 go 10 lá i ndiaidh an insteallta.
- Go hannamh, d'fhéadfadh páistí an 'leicneach bheag' a tholg le h-at sa ngiall sa tríú seachtain i ndiaidh an insteallta.

Níl siad seo tógálach.

D'fhéadfá paraicéiteamól nó iobúpróifein a thabhairt dod' pháiste chun faoiseamh a thabhairt ar phianta nó chun an fiabhras a íslíú.

Má tá imní ort faoi do pháiste, is féidir leat dul i dteagmháil leis an bhfoireann vacsaínithe scoile le linn uaireanta oifige, Luan go hAoine ar

Má theastaíonn comhairle leighis uait lasmuigh de na huaireanta seo, déan teagmháil le do Dhochtúir Teaghlaigh le do thoil. Tabharfar taifead de vacsaíní do pháiste duit nuair a bheidh siad ar fad faighte. **Coinnigh an taifead seo sábháilte.**

Tugadh na vacsaíní seo a leanas dod' pháiste inniu:

Tetravac ☐ MMRVaxPro ☐ Priorix ☐

05/2024
Cód Ordaithe: HNI00876

Tá tuilleadh eolais ar fáil ar: www.immunisation.ie

Chomh maith leis sin, gheobhaidh tú naisc leis na bileoga eolais d'othair le haghaidh na vacsaíní ag www.hpra.ie. Cuardaigh Tetravac (4 in 1), Priorix nó MMRVaxPro (MMR) chun an bhileog eolais chuí d'othair a léamh.