

## Treatment in community setting

Drug	Dose	Duration	+/- Notes
<b>Non-severe CDI</b>			
<ul style="list-style-type: none"> <li><b>Mildly symptomatic patients</b> (positive test result but less than 3 episodes of diarrhoea in 24 hours):</li> </ul> <p>It is recommended to discontinue antibiotic therapy with the inciting antibiotic if possible and closely monitor the patient for 48 hours. <i>C. difficile</i> treatment should be initiated if any signs of clinical deterioration are observed.</p>			
<ul style="list-style-type: none"> <li>Positive test result and <b>3 or more episodes of diarrhoea in 24 hours</b>, follow treatment options below:</li> </ul>			
<b>1<sup>st</sup> line option:</b>			
Metronidazole oral	400mg every 8 hours	10 days	
<b>2<sup>nd</sup> line option:</b>			
Vancomycin oral <sup>1</sup>	125mg every 6 hours	10 days	
<b>Severe CDI</b>			
<ul style="list-style-type: none"> <li>Refer to hospital</li> </ul>			
<b>First recurrence of CDI</b>			
<ul style="list-style-type: none"> <li>If non-severe CDI: see treatment recommendation below</li> <li>If severe CDI: refer to hospital</li> </ul>			
<b>1<sup>st</sup> line options:</b>			
Vancomycin oral <sup>1</sup>	125mg every 6 hours	10 days	Only use if metronidazole was used for treatment of the first episode
or			
Fidaxomicin oral	200mg every 12 hours	10 days	Discuss with a clinical microbiologist or infectious diseases consultant. High Tech item, see prescribing notes* If initial CDI episode was treated with fidaxomicin, seek micro/ID advice.

<sup>1</sup> Vancomycin: If a person has swallowing difficulties or a nasogastric or PEG tube for enteral administration, vials of vancomycin powder for injection may be used to make an extemporaneous oral solution. Vancomycin (oral) adult – includes information for dispensing pharmacists on how to make extemporaneous oral solution.

\*Fidaxomicin prescribing: Fidaxomicin should only be initiated on the recommendation of a Consultant Microbiologist or Infectious Diseases Physician. Fidaxomicin is only available from community pharmacies through the High Tech Arrangements. A GP may prescribe fidaxomicin, but must state the name and base hospital of the consulting Consultant Microbiologist or Infectious Diseases Physician on the prescription in order for the community pharmacy to process through the High Tech Arrangements. For hospital patients, follow the normal procedure for prescriptions of High Tech drugs.