

STI	Look back period	Empiric treatment	Window period
	Sexual partners should be informed of the need for testing and patients should be encouraged to inform their sexual partners		
<a href="#"><u>Chlamydia trachomatis</u></a>	Males with urethral symptoms: all contacts since symptom onset and in the 4 weeks prior to the symptoms. Asymptomatic Partner; Symptoms at all other sites (rectal, throat and eye): all contacts in prior 6 months	Yes	Two weeks
<b>Lymphogranuloma Venereum (LGV)</b>	Symptomatic: All contacts in the 4/52 prior to symptom onset Asymptomatic: all partners in past 3/12	N/A	
<a href="#"><u>Gonorrhoea (Neisseria Gonorrhoea)</u></a>	Males with urethral symptoms: all partners in past 2/52 or last partner if longer than 2/52 All others including asymptomatic infection: 3/12	Yes	Two weeks
<a href="#"><u>Trichomoniasis (Trichomonas Vaginalis)</u></a>	Sexual partners in the preceding 4 weeks prior to presentation	Yes	N/A
<b>HIV</b>	Refer to GUM/HIV clinic	Refer to GUM/HIV clinic (Be mindful of new sexual partners who may require PEP)	45 days
<a href="#"><u>Anogenital Warts (Human Papilloma Virus)</u></a>	N/A	N/A	N/A
<a href="#"><u>Syphilis (Treponema pallidum)</u></a>	Refer syphilis cases to GUM clinic.	Refer to GUM clinic	12 weeks
<b>Hepatitis</b>	Refer to GUM clinic	Refer to GUM clinic	12 weeks
<a href="#"><u>Genital Herpes (Herpes Simplex Virus)</u></a>	N/A	N/A	N/A
<a href="#"><u>Pelvic Inflammatory Disease</u></a>	Current male partners	N/A	
<a href="#"><u>Mycoplasma Genitalium</u></a>	Refer to GUM clinic	Refer to GUM clinic	As per GUM clinic