

## Darolutamide Therapy

### INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement Status
Treatment of adult men with non-metastatic castration-resistant prostate cancer (nmCRPC) who are at high risk of developing metastatic disease.	C61	00693a	CDS 01/03/2022

### TREATMENT:

*The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patient's individual clinical circumstances.*

Darolutamide is taken twice daily until disease progression or unacceptable toxicity develops.

Drug	Dose	Route	Cycle
Darolutamide	600mg twice daily	PO	Continuous
The tablets should be taken whole with food			
If a dose is missed, the dose should be taken as soon as the patient remembers prior to the next scheduled dose. The patient should not take two doses together to make up for a missed dose.			
Medical castration with a luteinising hormone-releasing hormone (LHRH) analogue should be continued during treatment of patients not surgically castrated.			

### ELIGIBILITY:

- Indications as above
- Age > 18
- Histologically or cytologically confirmed adenocarcinoma of prostate without neuroendocrine differentiation or small cell features.
- Diagnosis of castration-resistant prostate cancer
- Castrate level of serum testosterone (<1.7 nmol/L [50ng/dL]) on gonadotrophin releasing hormone agonist or antagonist therapy or after bilateral orchiectomy
- Baseline PSA level of at least 2 ng per millilitre
- PSA doubling time of 10 months or less
- ECOG 0-1

### CAUTION:

Use in caution in:

- Patients with clinically significant cardiovascular disease within past 6 months

NCCP Regimen: Darolutamide Therapy	Published: 09/02/2022 Review: 09/02/2023	Version number: 1
Tumour Group: Genitourinary NCCP Regimen Code: 00693	ISMO Contributor: Prof Maccon Keane	Page 1 of 4
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## EXCLUSIONS:

- Known hypersensitivity to darolutamide or any of the excipients .
- Patients with detectable metastases or a history of metastatic disease
- Prior treatment with second-generation androgen receptor antagonists
- Prior treatment with CYP17 inhibitors
- Prior chemotherapy or immunotherapy for prostate cancer, except adjuvant/neoadjuvant treatment
- Gastrointestinal disorder affecting absorption

## PRESCRIPTIVE AUTHORITY:

- The treatment plan must be initiated by a Consultant Medical Oncologist

## TESTS:

### Baseline tests:

- FBC, renal and liver profile
- Blood pressure
- ECG in patients at risk of QT prolongation

### Regular tests:

- FBC, renal and liver profile
- Blood pressure as clinically indicated
- ECG as clinically indicated

### Disease monitoring:

Disease monitoring should be in line with the patient's treatment plan and any other test/s as directed by the supervising Consultant.

## DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Consultant.

NCCP Regimen: Darolutamide Therapy	Published: 09/02/2022 Review: 09/02/2023	Version number: 1
Tumour Group: Genitourinary NCCP Regimen Code: 00693	ISMO Contributor: Prof Maccon Keane	Page 2 of 4
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a>  <i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## Renal and Hepatic Impairment:

**Table 1: Dose modification of Darolutamide in renal and hepatic impairment**

Renal Impairment	Hepatic Impairment
No dose adjustment is necessary for patients with mild or moderate renal impairment. For patients with severe renal impairment not receiving haemodialysis, the recommended starting dose is 300 mg twice daily.	No dose adjustment is necessary for patients with mild hepatic impairment. The available data on darolutamide pharmacokinetics in moderate hepatic impairment is limited. Darolutamide has not been studied in patients with severe hepatic impairment. For patients with moderate and severe hepatic impairment (Child-Pugh Classes B and C), the recommended starting dose is 300 mg twice daily

## Management of adverse events:

**Table 2: Dose Modification of Darolutamide for Adverse Events**

Adverse Reaction	Recommended Dose Modification
Intolerable or $\geq$ Grade 3 reaction	Withhold or reduce dose to 300mg twice daily until symptoms improve. Treatment may then be resumed at a dose of 600mg twice daily if tolerated. Dose reduction below 300mg twice daily is not recommended because efficacy has not been established

## SUPPORTIVE CARE:

**EMETOGENIC POTENTIAL:** Minimal (**Refer to local policy**).

**PREMEDICATIONS:** Not usually required

**OTHER SUPPORTIVE CARE:** None specified

## ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS:

*The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.*

- **Recent cardiovascular disease:** Patients with clinically significant cardiovascular disease in the past 6 months including stroke, myocardial infarction, severe/unstable angina pectoris, coronary/peripheral artery bypass graft, and symptomatic congestive heart failure were excluded from the clinical studies. Therefore, the safety of darolutamide in these patients has not been established. If darolutamide is prescribed, patients with clinically significant cardiovascular disease should be treated for these conditions according to established guidelines
- **Androgen deprivation therapy may prolong the QT interval:** In patients with a history of risk factors for QT prolongation and in patients receiving concomitant medicinal products that might prolong the QT

NCCP Regimen: Darolutamide Therapy	Published: 09/02/2022 Review: 09/02/2023	Version number: 1
Tumour Group: Genitourinary NCCP Regimen Code: 00693	ISMO Contributor: Prof Maccon Keane	Page 3 of 4
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

interval, physicians should assess the benefit-risk ratio including the potential for Torsade de pointes prior to initiating darolutamide.

- **Concomitant use with other medicinal products:** Use of strong CYP3A4 and P-gp inducers during treatment with darolutamide may decrease the plasma concentration of darolutamide and is not recommended, unless there is no therapeutic alternative. Selection of an alternate concomitant medicinal product with less potential to induce CYP3A4 or P-gp should be considered. Patients should be monitored for adverse reactions of BCRP, OATP1B1 and OATP1B3 substrates as co-administration with darolutamide may increase the plasma concentrations of these substrates. Co-administration with rosuvastatin should be avoided unless there is no therapeutic alternative.

## DRUG INTERACTIONS:

- Darolutamide is a substrate of CYP3A4 and P-glycoprotein (P-gp). Use of strong and moderate CYP3A4 inducers and P-gp inducers (e.g. carbamazepine, phenobarbital, St. John's Wort, phenytoin, and rifampicin) during treatment with darolutamide is not recommended, unless there is no therapeutic alternative.
- Concomitant use of darolutamide with a combined P-gp and strong CYP3A4 inhibitor increases darolutamide exposure which may increase the risk of darolutamide adverse reactions.
- Darolutamide is an inhibitor of breast cancer resistance protein (BCRP) and Organic Anion Transporting Polypeptides (OATP) 1B1 and 1B3. Co-administration of rosuvastatin should be avoided unless there is no therapeutic alternative. Selection of an alternative concomitant medicinal product with less potential to inhibit BCRP
- Since androgen deprivation treatment may prolong the QT interval, the co-administration with medicinal products known to prolong the QT interval or medicinal products able to induce Torsade de pointes should be carefully evaluated.

## REFERENCES:

1. Fizazi k, et al. Darolutamide in Non-metastatic, Castration-resistant Prostate Cancer. N Engl J Med 2019;380:1235-46
2. Darolutamide (Nubeqa®) Summary of Product Characteristics. Accessed January 2022 Available at: [https://www.ema.europa.eu/en/documents/product-information/nubeqa-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/nubeqa-epar-product-information_en.pdf)
3. NCCP Classification Document for Systemic Anti-Cancer Therapy (SACT) Induced Nausea and Vomiting. V3 2021. <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/nccp-classification-document-for-systemic-anti-cancer-therapy-sact-induced-nausea-and-vomiting.pdf>

Version	Date	Amendment	Approved By
1	09/02/2022		Prof Maccon Keane

Comments and feedback welcome at [oncologydrugs@cancercontrol.ie](mailto:oncologydrugs@cancercontrol.ie).

NCCP Regimen: Darolutamide Therapy	Published: 09/02/2022 Review: 09/02/2023	Version number: 1
Tumour Group: Genitourinary NCCP Regimen Code: 00693	ISMO Contributor: Prof Maccon Keane	Page 4 of 4
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a>  <i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		