

Avelumab Monotherapy

INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement Status
Treatment of adult patients with metastatic Merkel cell carcinoma who have received 1 or more lines of chemotherapy for metastatic disease	C4A	00535a	ODMS 01/05/2019

TREATMENT:

The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patients individual clinical circumstances.

Avelumab is administered on day 1 of a 14 day cycle until disease progression or unacceptable toxicity develops.

Day	Drug	Dose	Route	Diluent & Rate	Cycle
1	Avelumab	800mg	IV infusion	250mL NaCl 0.9% over 60 minutes	Every 14 days
Administer the solution for infusion using a sterile, non-pyrogenic, low-protein binding 0.2 micrometre in-line or add-on filter.					

ELIGIBILITY:

- Indication as above
- Histologically proven metastatic merkel cell carcinoma
- ECOG 0-1

EXCLUSIONS:

- Hypersensitivity to avelumab or any of the excipients
- Prior therapy with other immune checkpoint inhibitors
- Any medical condition that requires immunosuppressive doses of systemic corticosteroids or other immunosuppressive medication(s) (defined as >10mg prednisolone/daily (or steroid equivalent, excluding inhaled or topical steroids)
- Uncontrolled systemic infections such as HIV, Hepatitis B and Hepatitis C
- History of organ transplant
- Untreated, symptomatic CNS metastases

PRESCRIPTIVE AUTHORITY:

The treatment plan must be initiated by a Consultant Medical Oncologist

NCCP Regimen: Avelumab Monotherapy	Published: 01/05/2019 Review: 01/05/2021	Version number: 3
Tumour Group: Skin NCCP Regimen Code: 00535	ISMO Contributor: Dr Deirdre O'Mahony	Page 1 of 7
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

TESTS:

Baseline tests:

- FBC, renal and liver profile
- Blood glucose
- Thyroid function test
- Virology screen -Hepatitis B (HBsAg, HBcoreAb) & C, HIV.

Regular tests:

- FBC, renal and liver profile prior to each cycle
- Blood glucose and thyroid function test prior to each cycle

Disease monitoring:

Disease monitoring should be in line with the patient's treatment plan and any other test/s as directed by the supervising Consultant.

DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Consultant.
- Dose escalation or dose reduction is not recommended.
- Dosing delay or discontinuation may be required based on individual safety and tolerability

Renal and Hepatic Impairment:

Table 1: Dose modification of avelumab in renal and hepatic impairment

Renal Impairment	Hepatic Impairment
No dose adjustment is needed for patients with mild or moderate renal impairment. There are insufficient data in patients with severe renal impairment for dosing recommendations	No dose adjustment is needed for patients with mild hepatic impairment. There are insufficient data in patients with moderate or severe hepatic impairment for dosing recommendations

Management of adverse events:

Table 2: Dose Modification of avelumab for Adverse Events

Adverse reactions	Severity*	Recommended dose modification
Infusion-related reactions	Grade 1	Reduce infusion rate by 50%
	Grade 2	Withhold until adverse reactions recover to Grade 0-1; restart infusion with a 50% slower rate
	Grade 3 or Grade 4	Permanently discontinue
Pneumonitis	Grade 2	Withhold until adverse reactions recover to Grade 0-1
	Grade 3 or Grade 4 or recurrent Grade 2	Permanently discontinue

NCCP Regimen: Avelumab Monotherapy	Published: 01/05/2019 Review: 01/05/2021	Version number: 3
Tumour Group: Skin NCCP Regimen Code: 00535	ISMO Contributor: Dr Deirdre O'Mahony	Page 2 of 7
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

Hepatitis	Aspartate aminotransferase (AST) or alanine aminotransferase (ALT) greater than 3 and up to 5 times upper limit of normal (ULN) or total bilirubin greater than 1.5 and up to 3 times ULN	Withhold until adverse reactions recover to Grade 0-1
	AST or ALT greater than 5 times ULN or total bilirubin greater than 3 times ULN	Permanently discontinue
Colitis	Grade 2 or Grade 3 colitis or diarrhoea	Withhold until adverse reactions recover to Grade 0-1
	Grade 4 colitis or diarrhoea or recurrent Grade 3 colitis	Permanently discontinue
Pancreatitis	Suspected pancreatitis	Withhold
	Confirmed pancreatitis	Permanently discontinue
Myocarditis	Suspected myocarditis	Withhold
	Confirmed myocarditis	Permanently discontinue
Endocrinopathies (hypothyroidism, hyperthyroidism, adrenal insufficiency, hyperglycaemia)	Grade 3 or Grade 4	Withhold until adverse reactions recover to Grade 0-1
Nephritis and renal dysfunction	Serum creatinine more than 1.5 and up to 6 times ULN	Withhold until adverse reactions recover to Grade 0-1
	Serum creatinine more than 6 times ULN	Permanently discontinue
Other immune-related adverse reactions (including myositis, hypopituitarism, uveitis, Guillain-Barré syndrome)	For any of the following: • Grade 2 or Grade 3 clinical signs or symptoms of an immune-related adverse reaction not described above	Withhold until adverse reactions recover to Grade 0-1
	For any of the following: • Life threatening or Grade 4 adverse reaction (excluding endocrinopathies controlled with hormone replacement therapy) • Recurrent Grade 3 immune-related adverse reaction • Requirement for 10 mg per day or greater prednisone or equivalent for more than 12 weeks • Persistent Grade 2 or Grade 3 immune-mediate adverse reactions lasting 12 weeks or longer	Permanently discontinue

* Toxicity was graded per National Cancer Institute Common Terminology Criteria for Adverse Events Version 4.0 (NCI-CTCAE v4.03)

SUPPORTIVE CARE:

EMETOGENIC POTENTIAL: Minimal (**Refer to local policy**).

NCCP Regimen: Avelumab Monotherapy	Published: 01/05/2019 Review: 01/05/2021	Version number: 3
Tumour Group: Skin NCCP Regimen Code: 00535	ISMO Contributor: Dr Deirdre O'Mahony	Page 3 of 7
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

PREMEDICATIONS:

Patients have to be premedicated with an antihistamine and with paracetamol prior to the first 4 infusions of avelumab. If the fourth infusion is completed without an infusion-related reaction, premedication for subsequent doses should be administered at the discretion of the clinician

Table 4: Suggested pre-medications prior to avelumab infusion:

Drugs	Dose	Route
Paracetamol	1g	PO at least 30 minutes prior to avelumab infusion
Chlorphenamine	10mg	IV bolus at least 30 minutes prior to avelumab infusion

OTHER SUPPORTIVE CARE:

Women of childbearing potential should be advised to avoid becoming pregnant while receiving avelumab and should use effective contraception during treatment with avelumab and for at least 1 month after the last dose of avelumab.

ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS

The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.

- **Infusion-related reactions:** Infusion-related reactions, which might be severe, have been reported in patients receiving avelumab. Patients should be monitored for signs and symptoms of infusion-related reactions including pyrexia, chills, flushing, hypotension, dyspnoea, wheezing, back pain, abdominal pain, and urticaria. Guidelines for management of infusion –related reactions are in Table 2 above.
- **Immune-related adverse reactions:** Most immune-related adverse reactions with avelumab were reversible and managed with temporary or permanent discontinuation of avelumab, administration of corticosteroids and/or supportive care.
 - Based on the severity of the adverse reaction, avelumab should be withheld and corticosteroids administered.
 - If corticosteroids are used to treat an adverse reaction, a taper of at least 1 month duration should be initiated upon improvement.
 - In patients, whose immune-related adverse reactions cannot be controlled with corticosteroid use, administration of other systemic immunosuppressants may be considered.

Table 5: management of immune related adverse reactions

Adverse reaction	Withhold/ discontinue	Recommended action -1 st occurrence
Immune-related pneumonitis		
Patients should be monitored for signs and symptoms of immune-related pneumonitis and causes other than immune-related pneumonitis should be ruled out.		
Grade 2	Withhold until resolution	Suspected pneumonitis should be confirmed with radiographic imaging. Corticosteroids should be administered for Grade ≥ 2 events (initial dose of 1 to 2 mg/kg/day prednisone or equivalent, followed by a corticosteroid taper
Grade 3 or Grade 4 or recurrent	Permanently	

NCCP Regimen: Avelumab Monotherapy	Published: 01/05/2019 Review: 01/05/2021	Version number: 3
Tumour Group: Skin NCCP Regimen Code: 00535	ISMO Contributor: Dr Deirdre O'Mahony	Page 4 of 7
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

Grade 2	discontinue	
Immune-related colitis Patients should be monitored for signs and symptoms of immune-related colitis and causes other than immune-related colitis should be ruled out.		
Grade 2 or Grade 3	Withhold until resolution	Corticosteroids should be administered for Grade \geq 2 events (initial dose of 1 to 2 mg/kg/day prednisone or equivalent followed by a corticosteroid taper)
Grade 4 or recurrent Grade 3	Permanently discontinue	
Immune-related hepatitis Patients should be monitored for changes in liver function and symptoms of immune-related hepatitis and causes other than immune-related hepatitis should be ruled out.		
Grade 2	Withhold until resolution	Corticosteroids should be administered for Grade \geq 2 events (initial dose 1 to 2 mg/kg/day prednisone or equivalent, followed by a corticosteroid taper)
Grade 3 or Grade 4	Permanently discontinue	
Immune-related pancreatitis Patients should be monitored for signs and symptoms of immune-related pancreatitis. In symptomatic patients, obtain gastroenterology consultation and laboratory investigations (including imaging) to ensure the initiation of appropriate measures at an early stage.		
Suspected immune-related pancreatitis	Withhold until resolution	Corticosteroids should be administered for immune-related pancreatitis (initial dose of 1 to 2 mg/kg/day prednisone or equivalent followed by a corticosteroid taper).
Confirmed immune-related pancreatitis	Permanently discontinue	
Immune-related myocarditis Patients should be monitored for signs and symptoms of immune-related myocarditis. In symptomatic patients, obtain cardiologic consultation and laboratory investigations to ensure the initiation of appropriate measures at an early stage.		
Suspected immune-related myocarditis	Withhold until resolution	Corticosteroids should be administered for immune-related myocarditis (initial dose of 1 to 2 mg/kg/day prednisone or equivalent followed by a corticosteroid taper). If no improvement within 24 hours on corticosteroids, additional immunosuppression (e.g., mycophenolate, infliximab, anti-thymocyte globulin) should be considered.
Confirmed immune-related myocarditis	Permanently discontinue	
Immune-related endocrinopathies Immune-related thyroid disorders, immune-related adrenal insufficiency, and Type 1 diabetes mellitus have been reported in patients receiving avelumab. Patients should be monitored for clinical signs and symptoms of endocrinopathies.		
Thyroid disorders: (<i>hypothyroidism/hyperthyroidism</i>)		Hypothyroidism should be managed with replacement therapy and hyperthyroidism with anti-thyroid medicinal product as needed
Grade 3 or 4	Withhold until	

NCCP Regimen: Avelumab Monotherapy	Published: 01/05/2019 Review: 01/05/2021	Version number: 3
Tumour Group: Skin NCCP Regimen Code: 00535	ISMO Contributor: Dr Deirdre O'Mahony	Page 5 of 7
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

	resolution	
Adrenal insufficiency Grade 3 or Grade 4 symptomatic	Withhold	Corticosteroids should be administered (1 to 2 mg/kg/day prednisone intravenously or oral equivalent) for Grade ≥ 3 adrenal insufficiency followed by a taper until a dose of less than or equal to 10 mg/day has been reached.
Type 1 diabetes mellitus Grade ≥ 3 hyperglycaemia	Withhold	Avelumab can cause Type 1 diabetes mellitus, including diabetic ketoacidosis. Patients should be monitored for hyperglycaemia or other signs and symptoms of diabetes. Initiate treatment with insulin for Type 1 diabetes mellitus Antihyperglycaemics should be administered. Treatment with avelumab should be resumed when metabolic control is achieved on insulin replacement therapy
Immune-related nephritis and renal dysfunction Grade 2 or 3 nephritis Grade 4	Withhold until resolution to ≤ Grade 1 Permanently discontinue	Corticosteroids (initial dose of 1 to 2 mg/kg/day prednisone or equivalent followed by a corticosteroid taper) should be administered for Grade ≥ 2 nephritis..
Other immune-related adverse reactions Other clinically important immune-related adverse reactions were reported in less than 1% of patients: myocarditis including fatal cases, myositis, pancreatitis, hypopituitarism, uveitis, and Guillain-Barré syndrome		

DRUG INTERACTIONS:

- No interaction studies have been conducted with avelumab.
- Avelumab is primarily metabolised through catabolic pathways, therefore, it is not expected that avelumab will have pharmacokinetic drug-drug interactions with other medicinal products.
- Current drug interaction databases should be consulted for more information.

ATC CODE:

Avelumab - L01XC31.

REFERENCES:

1. Kaufman et al. Updated efficacy of avelumab in patients with previously treated metastatic Merkel cell carcinoma after ≥1 year of follow-up: JAVELIN Merkel 200, a phase 2 clinical trial. Journal for ImmunoTherapy of Cancer (2018) 6:7
2. Kaufman et al. Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial Lancet Oncol 2016; 17: 1374–85
3. Avelumab (Bavencio®) 20 mg/mL concentrate for solution for infusion Summary of Product characteristics. Last updated 04/11/2019. Accessed Dec 2019 available at https://www.ema.europa.eu/en/documents/product-information/bavencio-epar-product-information_en.pdf

NCCP Regimen: Avelumab Monotherapy	Published: 01/05/2019 Review: 01/05/2021	Version number: 3
Tumour Group: Skin NCCP Regimen Code: 00535	ISMO Contributor: Dr Deirdre O'Mahony	Page 6 of 7
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

Version	Date	Amendment	Approved By
1	10/04/2019		Dr Deirdre O'Mahony
2	09/07/2019	Updated immune related adverse reactions to include pancreatitis as per SPC update	Dr Deirdre O'Mahony
3	14/02/2020	Updated dosing posology as per SmPC update to flat dosing. Updated dose modification and adverse events for pancreatitis and myocarditis as per SmPC update	Dr Deirdre O'Mahony

Comments and feedback welcome at oncologydrugs@cancercontrol.ie.

NCCP Regimen: Avelumab Monotherapy	Published: 01/05/2019 Review: 01/05/2021	Version number: 3
Tumour Group: Skin NCCP Regimen Code: 00535	ISMO Contributor: Dr Deirdre O'Mahony	Page 7 of 7
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		