

## Cyclophosphamide/Total Body Irradiation (TBI)–MAC–MUD

### INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement Status
Myeloablative conditioning (MAC) for matched unrelated donor allogeneic stem cell transplant in patients with lymphoid disorders and acute or chronic leukaemias.	C91	00631a	Hospital

### TREATMENT:

Conditioning chemotherapy is administered over **2 days**. Stem cells are infused on **day 0**.

Facilities to treat anaphylaxis **MUST** be present when conditioning therapy and stem cells are administered.

Day (time)	Drug	Dose	Route	Diluent & Rate
-5, -4 (09.30)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-5, -4 (10.00)*	Cyclophosphamide	60mg/kg	IV infusion	1000ml sodium chloride 0.9% over 3 hours
-5, -4 (13.00)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-5, -4 (16.00)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-5, -4 (19.00)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-5, -4 (22.00)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-4, -3 (02.00)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-4, -3 (06.00)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-3 (10.00)*	Mesna	24mg/kg	Slow IV push	Into side arm of fast flowing sodium chloride 0.9% infusion
-3,-2,-1	Fractionated TBI	Twice Daily	n/a	n/a
0	Stem cell infusion			
+1 (at least 24 hours post completion of stem cell infusion)	Methotrexate <sup>a</sup>	15mg/m <sup>2</sup>	IV infusion	50ml sodium chloride 0.9% over 10 minutes
+3, +6, +11	Methotrexate	10mg/m <sup>2</sup>	IV infusion	50ml sodium chloride 0.9% over 10 minutes
<b>Dose rounding:</b> Mesna to the nearest 100mg, Cyclophosphamide to the nearest 20mg, Methotrexate to the nearest 2.5mg				
<sup>a</sup> Day +1 methotrexate should be administered at least 24 hours post completion of stem cell infusion. In the event where this timing results in methotrexate being infused during the night, it is reasonable to reschedule the administration time of the day +3 methotrexate dose to the next morning, to avoid administration during the night. The amended administration timing can then be maintained for subsequent methotrexate doses.				
*Denotes recommended administration times				

NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD	Published: 06/08/2021 Review: 01/04/2025	Version number: 1a
Tumour Group: Transplant NCCP Regimen Code: 00631	IHS Contributor: SJH Stem Cell Transplant Group	Page 1 of 8
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## ELIGIBILITY:

- Indications as above
- Medical assessment as per SJH BMT assessment form

## EXCLUSIONS:

- Hypersensitivity to cyclophosphamide, mesna, methotrexate or any of the excipients
- Pregnancy and lactation

## PRESCRIPTIVE AUTHORITY:

- The treatment plan must be initiated by a Haematology Consultant working in the area of stem cell transplantation in a unit suitable for carrying out this treatment.

## TESTS:

- Baseline and regular tests in accordance with SJH Haematopoietic Stem Cell Transplant work-up protocols

### Disease monitoring:

Disease monitoring should be in line with the patient's treatment plan and any other test/s as directed by the supervising Consultant.

NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD	Published: 06/08/2021 Review: 01/04/2025	Version number: 1a
Tumour Group: Transplant NCCP Regimen Code: 00631	IHS Contributor: SJH Stem Cell Transplant Group	Page 2 of 8
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Haematology Consultant.
- **Chemotherapy dosing in obese adult patients:** For patients with a BMI > 30kg/m<sup>2</sup> please refer to 'Chemotherapy Dosing in Obese Adult Stem Cell Transplant Recipients – Guidelines' for guidance on individual drug dosing as per SJH policy available on the SJH intranet.
- **Renal and Hepatic Impairment:**
  - Dose modifications are generally not undertaken in conditioning regimens.
  - Discuss with the consultant if the creatinine clearance is < 50 ml/min or if abnormal hepatic function.
  - Consult the following resources to inform any renal or hepatic dose modification discussions:
    - Summary of product characteristics (SPC) available at <http://www.hpra.ie>
    - Krens et al Lancet Oncol 2019;20(4) e200-e207 "Dose Recommendations for anticancer drugs in patients with renal or hepatic impairment" available at <https://pubmed.ncbi.nlm.nih.gov/30942181/>
    - UCHL renal impairment guidelines and hepatic impairment guidelines available on SJH intranet

## SUPPORTIVE CARE:

### Antiemetics:

Table 1: Recommended SJH Regimen Specific Antiemetics

Prevention of acute nausea and vomiting			Prevention of delayed nausea and vomiting			Comment
Drug	Dose	Admin Day	Drug	Dose	Admin Day	
Dexamethasone	12mg PO	-5, -4	Dexamethasone	8mg PO	-3, -2, -1	Exclude aprepitant due to interaction with cyclophosphamide
Ondansetron	8mg PO/IV TDS	-5, -4				

### Cyclophosphamide hydration and diuresis:

- Pre stem cell infusion: Start pre-hydration at 6.00 am on Day -5
  - Recommended hydration regimen is sodium chloride 0.9% 2-3L/m<sup>2</sup> over 24 hours
- Continue hydration for at least 24 hours after completion of cyclophosphamide
- Diuretics may be indicated for positive fluid balance, weight gain or declining urine production (<100ml/m<sup>2</sup>/hr)
  - Furosemide 20-40mg IV PRN should be prescribed

NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD	Published: 06/08/2021 Review: 01/04/2025	Version number: 1a
Tumour Group: Transplant NCCP Regimen Code: 00631	IHS Contributor: SJH Stem Cell Transplant Group	Page 3 of 8
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## Other Supportive Care:

**Table 2: Other Supportive Medication**

<p><b>GvHD prophylaxis</b> Refer to signed off BMT assessment form for confirmed <b>choice and target level</b> of immunosuppression</p>	<p><b>Tacrolimus</b> Tacrolimus 0.03mg/kg once daily IV over 22 hours from day -1</p> <ul style="list-style-type: none"> <li>The equivalent oral dose is: (Total IV dose) twice daily PO</li> <li>Target levels: 5-10 nanograms/ml</li> </ul>
<p><b>GvHD and VOD prophylaxis</b></p>	<ul style="list-style-type: none"> <li>Ursodeoxycholic acid 250mg TDS PO</li> <li>Continue until day +90</li> </ul>
<p><b>HSV prophylaxis</b></p>	<p>All patients should receive the following until CD4 count &gt;200/microlitre:</p> <ul style="list-style-type: none"> <li>Valaciclovir 500mg once daily PO</li> </ul> <p><b>or</b></p> <ul style="list-style-type: none"> <li>Aciclovir 250mg TDS IV (if oral route not available or ANC &lt; 0.5X10<sup>9</sup>/L)</li> </ul> <p>Patients with an active herpes infection should receive the following:</p> <ul style="list-style-type: none"> <li>Valaciclovir 1g TDS PO</li> </ul> <p><b>or</b></p> <ul style="list-style-type: none"> <li>Aciclovir 10mg/kg TDS IV (if oral route not available)</li> </ul>
<p><b>CMV prophylaxis</b>  Prescribe for all CMV seropositive recipients</p>	<p><b>Patients receiving CMV prophylaxis with letermovir also require HSV prophylaxis above</b></p> <ul style="list-style-type: none"> <li>Letermovir 480mg once daily PO/IV, as appropriate, starting Day +1 if patient is receiving tacrolimus immunosuppression</li> <li>Letermovir via the oral route is first line.</li> <li>Letermovir IV at the same oral dose should be prescribed only where the patient cannot tolerate oral or where there are concerns around absorption.</li> <li>CMV prophylaxis is usually continued until day +100</li> </ul> <p>Patients should bring their oral letermovir supply with them on admission. High tech prescription will have been provided to patient at their counselling appointment pre-admission. Liaise with transplant pharmacist if any supply issues arise.</p> <p>When ANC &gt; 1.0 x 10<sup>9</sup>/L, pre-emptive monitoring (9mls in EDTA [purple tube] (Tuesday and Fridays) should be carried out for CMV reactivation/infection in <u>all</u> patients</p>

<p>NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD</p>	<p>Published: 06/08/2021 Review: 01/04/2025</p>	<p>Version number: 1a</p>
<p>Tumour Group: Transplant NCCP Regimen Code: 00631</p>	<p>IHS Contributor: SJH Stem Cell Transplant Group</p>	<p>Page 4 of 8</p>
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a> <i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

<p><b>Antifungal prophylaxis</b> Refer to signed off BMT assessment form for confirmed choice of antifungal prophylaxis</p>	<p>When ANC <math>&lt;0.5 \times 10^9/L</math> or if patients on high dose steroids:</p> <ul style="list-style-type: none"> <li>Liposomal amphotericin 1mg/kg once daily IV Mon/Wed/Fri <b>or</b></li> <li>Caspofungin 70mg once daily IV Mon/Wed/Fri</li> </ul> <p>If at higher risk due to prior possible/probable fungal infection:</p> <ul style="list-style-type: none"> <li>Liposomal amphotericin 1mg/kg once daily IV <b>or</b></li> <li>Caspofungin 70mg once daily IV if <math>&gt;80kg</math> <b>or</b></li> <li>Caspofungin 70mg once daily IV on day 1 of treatment followed by 50mg once daily IV thereafter if <math>&lt;80kg</math></li> </ul>
<p><b>PJP prophylaxis</b></p>	<p><u>1st line therapy:</u></p> <ul style="list-style-type: none"> <li>Co-trimoxazole 960mg BD Mon/Wed/Fri PO</li> <li>Commence only on engraftment when ANC <math>&gt; 1.0 \times 10^9/L</math> if appropriate</li> </ul> <p><u>2nd line therapy (if allergic to co-trimoxazole or contraindicated):</u> <i>PJP Prophylaxis and T. gondii IgG NEGATIVE:</i></p> <ul style="list-style-type: none"> <li>Pentamidine 300mg nebule and salbutamol 2.5mg nebule pre-pentamidine, every 4 weeks <b>plus</b></li> <li>Phenoxyethylpenicillin 333mg BD daily PO</li> </ul> <p>Continue the phenoxyethylpenicillin until patients have been revaccinated and have adequate pneumococcal/haemophilus titres</p> <p><i>PJP Prophylaxis and T gondii IgG POSITIVE:</i></p> <ul style="list-style-type: none"> <li>Atovaquone 750mg BD PO plus</li> <li>Pyrimethamine 25mg once daily PO plus</li> <li>Folinic acid 15mg once daily PO plus</li> <li>Phenoxyethylpenicillin 333mg BD daily PO</li> </ul> <p>Continue the phenoxyethylpenicillin until patients have been revaccinated and have adequate pneumococcal/haemophilus titres</p> <p>Please note: If a patient is to be discharged on atovaquone, pyrimethamine or folinic acid, please contact pharmacy in advance to arrange supply and funding through a community drugs scheme</p>
<p><b>Mouthcare</b></p>	<p>Mucositis WHO grade <math>&lt; 2</math>:</p> <ul style="list-style-type: none"> <li>Sodium chloride 0.9% 10ml QDS mouthwash</li> <li>Nystatin 1ml QDS PO (use 15 minutes after sodium chloride 0.9% mouthwash)</li> </ul> <p>Mucositis WHO grade <math>\geq 2</math>:</p> <ul style="list-style-type: none"> <li>Chlorhexidine digluconate 0.12% (Kin<sup>®</sup> mouthwash) 10mls QDS mouthwash</li> <li>Nystatin 1ml QDS PO (use 15 minutes after Kin<sup>®</sup> mouthwash)</li> </ul>

<p>NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD</p>	<p>Published: 06/08/2021 Review: 01/04/2025</p>	<p>Version number: 1a</p>
<p>Tumour Group: Transplant NCCP Regimen Code: 00631</p>	<p>IHS Contributor: SJH Stem Cell Transplant Group</p>	<p>Page 5 of 8</p>
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a> This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></p>		

<b>Gastroprotection</b>	<ul style="list-style-type: none"> <li>Lansoprazole 30mg / omeprazole 40mg once daily PO <b>or</b></li> <li>Esomeprazole 40mg once daily IV (if oral route not available)</li> </ul>
<b>Folate supplementation</b>	<p><b>Methotrexate is included as GvHD prophylaxis. Folinic acid should not be administered on the same days as methotrexate.</b></p> <p>The first dose of folinic acid must be administered at a minimum of 24 hours post completion of methotrexate. Prescribe as outlined below:</p> <ul style="list-style-type: none"> <li>Folinic acid 15mg once daily IV on days <b>+2,+4,+5,+7,+8,+9,+10 and +12 onwards</b></li> <li>Switch to folic acid 5mg once daily PO when oral route is available</li> </ul>
<b>Vitamin K supplementation</b>	<p>Beginning on day +2 post stem cell transplant</p> <ul style="list-style-type: none"> <li>Vitamin K (phytomenadione) 10mg once weekly IV</li> </ul>
<b>Prevention of vaginal bleeding</b>	<p>If required for menstruating female patients until platelets &gt; 50 x10<sup>9</sup>/L</p> <ul style="list-style-type: none"> <li>Norethisterone 5mg TDS PO if &gt;55kg</li> <li>Norethisterone 5mg BD PO if &lt;55kg</li> </ul>
<b>Tumour Lysis syndrome</b>	<p>Consider allopurinol in active disease pre transplant</p> <ul style="list-style-type: none"> <li>Allopurinol 300mg once daily PO for 5-7 days and review</li> </ul>
<b>Hepatitis B prophylaxis/treatment</b>	<p>A virology screen is completed as part of transplant workup. Hepatitis B prophylaxis or treatment may be initiated in consultation with a Virology Consultant or Hepatology Consultant if required.</p> <p>Options may include:</p> <ul style="list-style-type: none"> <li>Lamivudine 100mg once daily PO <b>or</b></li> <li>Entecavir 500mcg once daily PO</li> </ul>
<b>Prevention of constipation</b>	<p>Consider laxatives if appropriate e.g.</p> <ul style="list-style-type: none"> <li>Senna two tablets (15mg) nocte PO while on ondansetron</li> </ul>
<b>Antibiotic standing order</b>	<p>Antibiotic standing order should be prescribed for neutropenic sepsis/neutropenic fever based on previous microbiology and renal function</p> <ul style="list-style-type: none"> <li>Piptazobactam 4.5g QDS IV <b>plus</b></li> <li>Amikacin* 15mg/kg once daily IV</li> </ul> <p>*Ciprofloxacin 400mg BD IV may be considered instead of amikacin in cases of renal impairment</p> <p>Refer to Antimicrobial Guidelines in the Prescriber's Capsule for antibiotic choice where a patient is allergic to any of the above</p>
<b>Magnesium and potassium standing order</b>	<p>Magnesium and potassium standing orders should be prescribed for all transplant patients in accordance with stem cell unit practice as indicated on EPMAR</p>
<b>VTE prophylaxis</b>	<p>Consider VTE prophylaxis in accordance with SJH policy</p>
<b>Bone Health</b>	<p>Consider calcium and vitamin D supplementation prior to discharge for patients who are on high dose steroids. Other medications for maintenance of bone health may need to be considered as appropriate.</p> <ul style="list-style-type: none"> <li>Calcium carbonate and colecalciferol (Caltrate® 600mg/400unit) one tablet BD</li> </ul>

NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD	Published: 06/08/2021 Review: 01/04/2025	Version number: 1a
Tumour Group: Transplant NCCP Regimen Code: 00631	IHS Contributor: SJH Stem Cell Transplant Group	Page 6 of 8
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS:

- Please refer to the relevant Summary of Product Characteristics and SJH Stem Cell Transplant Programme PPGs for full details.

## DRUG INTERACTIONS:

- The relevant Summary of Product Characteristics and current drug interaction databases should be consulted.

## REFERENCES:

1. Socie G, Clift RA, Blaise D, et al. Busulfan plus cyclophosphamide compared with total body irradiation plus cyclophosphamide before marrow transplantation for myeloid leukaemia: long term follow up of 4 randomised studies: The American Society of Haematology 2001; 98 (13):3569-73
2. Improved survival with ursodeoxycholic acid prophylaxis in allogenic stem cell transplantation: Long-term follow-up of a randomised study. *Biology of Blood and Marrow Transplantation* 2014; 20(1):135-138. Available at <https://pubmed.ncbi.nlm.nih.gov/24141008/>
3. Venous-occlusive disease/sinusoidal obstruction syndrome after haematopoietic stem cell transplantation: Middle East/North Africa regional consensus on prevention, diagnosis and management. *Bone Marrow Transplantation* 2017 Apr;52(4):588-591. Available at <https://pubmed.ncbi.nlm.nih.gov/27892944/>
4. Krens S D, Lassche, Jansman G F G A, et al. Dose recommendations for anticancer drugs in patients with renal or hepatic impairment. *Lancet Onco*/2019; 20:e201-08. [https://doi.org/10.1016/S1470-2045\(19\)30145-7](https://doi.org/10.1016/S1470-2045(19)30145-7)
5. Dosage Adjustment for Cytotoxics in Renal Impairment January 2009; North London Cancer Network.
6. Dosage Adjustment for Cytotoxics in Hepatic Impairment January 2009; North London Cancer Network.
7. NCCP Classification Document for Systemic Anti-Cancer Therapy (SACT) Induced Nausea and Vomiting. V3 2021. Available at: <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/nccp-classification-document-for-systemic-anti-cancer-therapy-sact-induced-nausea-and-vomiting.pdf>
8. Endoxana Injection 500 mg Powder for Solution for Injection. Summary of Product Characteristics. Accessed Nov 2020. Available at: [https://www.hpra.ie/img/uploaded/swedocuments/Licence\\_PA2299-027-001\\_21122018112107.pdf](https://www.hpra.ie/img/uploaded/swedocuments/Licence_PA2299-027-001_21122018112107.pdf)
9. Uromitexan 100mg/ml Solution for Injection or Infusion. Summary of Product Characteristics. Accessed Nov 2020. Available at: [https://www.hpra.ie/img/uploaded/swedocuments/Licence\\_PA2299-024-001\\_22102019104556.pdf](https://www.hpra.ie/img/uploaded/swedocuments/Licence_PA2299-024-001_22102019104556.pdf)
10. Methotrexate 1 g/10 ml Injection. Summary of Product Characteristics. Accessed Nov 2020.

NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD	Published: 06/08/2021 Review: 01/04/2025	Version number: 1a
Tumour Group: Transplant NCCP Regimen Code: 00631	IHS Contributor: SJH Stem Cell Transplant Group	Page 7 of 8
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

Available at: [https://www.hpra.ie/img/uploaded/swedocuments/Licence\\_PA0822-206-006\\_19052021104201.pdf](https://www.hpra.ie/img/uploaded/swedocuments/Licence_PA0822-206-006_19052021104201.pdf)

Version	Date	Amendment	Approved By
1	06/08/2021		SJH Stem Cell Transplant Group
1a	09/07/2024	Extension of review date as agreed with clinical reviewer	NCCP

Comments and feedback welcome at [oncologydrugs@cancercontrol.ie](mailto:oncologydrugs@cancercontrol.ie).

NCCP Regimen: Cyclophosphamide/Total Body Irradiation(TBI)–MAC–MUD	Published: 06/08/2021 Review: 01/04/2025	Version number: 1a
Tumour Group: Transplant NCCP Regimen Code: 00631	IHS Contributor: SJH Stem Cell Transplant Group	Page 8 of 8
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		