



Background: In recent years around 1,800 people are diagnosed with lung cancer annually – approximately 1,100 men and 700 women. Incidence rates are decreasing for men but increasing for women. Less than 1% of all new cases occur before the age of 40. More than 90% of patients are symptomatic at presentation. Currently over a third of patients have distant metastases by the time of diagnosis.

Risk Factors: Smoking, including passive smoking and smoking marijuana; radon exposure; exposure to heavy metals such as arsenic; radiation; asbestos dust; previous history of cancer (e.g. head & neck cancer). Smoking avoidance/cessation is the most important preventive strategy as over 90% of lung cancer can be attributed to cigarette smoke. However lung cancer can occur in patients without any of the listed risk factors.

ABOUT RAPID ACCESS SERVICES FOR LUNG CANCER

Rapid Access Services provide initial investigations, such as CT and bronchoscopy, to patients with suspected lung cancer, usually within one or two hospital visits. This reduces multiple hospital visits and patient anxiety and shortens the time period to diagnosis.

Patients should be assessed in the lung cancer rapid access service by a respiratory physician within **2 weeks** of receipt of request.

Who can refer your patient to the rapid access service?

- you, the GP
- a radiologist (in conjunction with GP)
- another hospital based clinician, e.g. from the Emergency Department

Who should you refer to the Rapid Access Service?

• a patient whose chest x-ray is suspicious of lung cancer.

Please include details on the referral of the hospital in which the investigation was carried out. If this was a different hospital / clinic, please fax or post a copy of the result to the clinic and request a copy of the film for the patient to bring with them if possible.

• a patient who has **haemoptysis**, or other symptoms which are concerning or persistent, even if their chest x-ray is normal.

When is referral to the rapid access service not appropriate?

If a patient presents with life threatening symptoms, an emergency referral should be made in the usual manner.

Table 1: Indications for Urgent Chest X-ray¹

A patient with the following signs or symptoms should be referred for urgent chest x-ray. A report should be back to the GP within one week of request.

Symptoms

- Haemoptysis
- New onset unexplained or persistent cough (>3 weeks)
- Alteration in character/severity of chronic cough
- Unexplained chest pain or dyspnoea
- Unexplained weight loss/cachexia
- Unexplained bone pain/neurological symptoms

Signs


- Clubbing
- Lymphadenopathy
- Focal chest signs
- Hepatomegaly

Note: if there is a suspicion of lung cancer, it is not advisable to delay referral by ordering an outpatient CT. A rapid access service can arrange both imaging and bronchoscopy.

1. Irish Thoracic Society Lung Cancer Sub-committee. Guidelines for the diagnosis and treatment of Lung Cancer. 3rd ed. Irish Thoracic Society 2009.

This guideline represents the view of the NCCP, which was arrived at after careful consideration of the evidence available. Health professionals are expected to take it fully into account when exercising their clinical judgement. The guidance does not, however, override the individual responsibility of health professionals to make decisions appropriate to each patient. This guideline will be reviewed as new evidence emerges, and supersedes all previous HSE/NCCP lung cancer GP referral guidelines.

REFERRAL PATHWAY



Notes: In certain hospitals, radiology may trigger referral to a rapid access service but this should not be assumed. In general, you (the GP) will be asked to inform the patient that they are being referred to the service.

GENERAL RECOMMENDATIONS

A patient who presents with symptoms suggestive of lung cancer should be referred to a team specialising in the management of lung cancer. To make a referral, send a National Lung Cancer Rapid Access Service Referral Form or submit an electronic referral form via Healthlink

(www.healthlink.ie). Additional referral forms can be obtained by phoning the National Cancer Control Programme at (01) 8287100 or logging onto www.cancercontrol.hse.ie

GPs should refer any patient they consider to have a high likelihood of lung cancer.

LUNG CANCER RAPID ACCESS SERVICES

Beaumont Hospital, PO Box 1297, Dublin 9	Tel: (01) 809 3484	
Cork University Hospital, Wilton, Cork	Tel: (021) 492 0453	Fax: (021) 492 2391
Galway University Hospital	Tel: (091) 542 234	Fax: (091) 542 092
Mater University Hospital, Dublin 7	Tel: (01) 803 2644/2295	Fax: (01) 803 4036
Mid Western Regional Hospital, Limerick	Tel: (061) 585 637	Fax: (061) 482 572
St. James's Hospital, Dublin 8	Tel: (01) 416 2196	Fax: (01) 410 3549
St. Vincent's University Hospital, Dublin 4	Tel: (01) 221 3702	Fax: (01) 221 3576
Waterford Regional Hospital, Waterford	Tel: (051) 848 988	Fax: (051) 848 844