



After your **COVID-19 Vaccine Janssen**

Thank you for protecting yourself and others by getting the vaccine. Now that you have had your vaccine, we ask you to read this document carefully, so you know how you can expect to feel in the next few days and where to get more information. Do read the COVID-19 vaccine information leaflet we gave you about the vaccine too.

We are also giving you a record of your vaccination today.

Please keep the record card safe.

What might happen in the next few days?

Some people who got the vaccine that you got today will:

- have swelling, redness and/or pain in the arm where they had the vaccine injection
- feel tired
- get a headache
- have a cough
- have muscle pain
- have joint pain
- have nausea
- get a fever – temperature of 38 degrees Celsius or above.

There have been very rare cases of unusual blood clots with low platelets in the blood in the first few weeks after vaccination. Platelets help the blood to clot.

You should seek prompt medical assistance and mention your recent vaccination if you:

- are breathless
- have pain in the chest or stomach
- have swelling or coldness in a leg
- get a severe or worsening headache (particularly three or more days after the vaccine)
- have blurred vision, confusion or seizures
- have persistent bleeding under the skin where there was no previous injury
- have many small bruises, reddish or purplish spots, or blood blisters under the skin.

There have been a very small number of cases of capillary leak syndrome in the first few days after vaccination. You should seek prompt medical assistance and mention your recent vaccination if you:

- have rapid swelling of the limbs – arms and legs
- have sudden weight gain

You may also feel faint due to low blood pressure.

There have been a very small number of cases of Guillain-Barré Syndrome after this vaccine.

Get urgent medical attention if you:

- have double vision or difficulty moving your eyes
- have difficulty swallowing, speaking or chewing
- have coordination problems, feel unsteady or have difficulty walking
- feel weakness in your arms, legs, chest or face
- have tingling in your hands or feet
- have problems with bladder control and bowel function

As with all vaccines, you can report suspected side effects to the Health Products Regulatory Authority HPRA. To report side effects to a COVID-19 vaccine, please visit

www.hpra.ie/report

What if I have a fever or have aches and pains?

If you have a fever or you have aches and pains, you can take paracetamol or ibuprofen to help. However, if your fever lasts more than 48 hours or if you are still concerned, please seek medical advice.

After vaccination

Getting a COVID-19 vaccine should protect you from the serious complications of COVID-19. It takes 14 days after getting your vaccine for it to work.

There is a chance you might still get COVID-19 even if you have the vaccine.

We do not know yet if having the vaccine stops you spreading the virus to others, so you must continue to follow public health advice:

- follow social distancing guidelines – keep two metres apart from others where possible
- wear a face covering
- wash your hands regularly.

Have a question?

If you are unsure about anything, or have any questions about the COVID-19 vaccine please ask your vaccinator today.

Visit www.hse.ie/covid19vaccine for more information on the COVID-19 vaccine.

Side Effects

Please report any side effects to the HPRA at
www.hpra.ie/report

Large print copies

This is a large print leaflet. The leaflet will be updated as new information becomes available. For a printed copy of the most recent version, contact NCBI by emailing library@ncbi.ie

Published by HSE on 29 July 2021.

Version 5.



Rialtas na hÉireann
Government of Ireland